

(12) INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(19) World Intellectual Property Organization
International Bureau



PCT



(43) International Publication Date
10 January 2008 (10.01.2008)

(10) International Publication Number
WO 2008/005562 A2

(51) International Patent Classification:
A61K 48/00 (2006.01)

(21) International Application Number:
PCT/US2007/015638

(22) International Filing Date: 9 July 2007 (09.07.2007)

(25) Filing Language: English

(26) Publication Language: English

(30) Priority Data:
60/819,704 7 July 2006 (07.07.2006) US

(71) Applicant (for all designated States except US): **UNIVERSITY OF MASSACHUSETTS** [US/US]; 225 Franklin Street, 12th Floor, Boston, MA 02110 (US).

(72) Inventors; and

(75) Inventors/Applicants (for US only): **ARONIN, Neil** [US/US]; 19 Whittier Road, Newtonville, MA 02460 (US). **ZAMORE, Phillip, D.** [US/US]; 500 Green Street, Northboro, MA 01532 (US).

(74) Agents: **MANDRAGOURAS, Amy, E.** et al.; Lahive & Cockfield, LLP, One Post Office Square, Boston, MA 02109-2127 (US).

(81) Designated States (unless otherwise indicated, for every kind of national protection available): AE, AG, AL, AM, AT, AU, AZ, BA, BB, BG, BH, BR, BW, BY, BZ, CA, CH, CN, CO, CR, CU, CZ, DE, DK, DM, DO, DZ, EC, EE, EG, ES, FI, GB, GD, GE, GH, GM, GT, HN, HR, HU, ID, IL, IN, IS, JP, KE, KG, KM, KN, KP, KR, KZ, LA, LC, LK, LR, LS, LT, LU, LY, MA, MD, ME, MG, MK, MN, MW, MX, MY, MZ, NA, NG, NI, NO, NZ, OM, PG, PH, PL, PT, RO, RS, RU, SC, SD, SE, SG, SK, SL, SM, SV, SY, TJ, TM, TN, TR, TT, TZ, UA, UG, US, UZ, VC, VN, ZA, ZM, ZW.

(84) Designated States (unless otherwise indicated, for every kind of regional protection available): ARIPO (BW, GH, GM, KE, LS, MW, MZ, NA, SD, SL, SZ, TZ, UG, ZM, ZW), Eurasian (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European (AT, BE, BG, CH, CY, CZ, DE, DK, EE, ES, FI, FR, GB, GR, HU, IE, IS, IT, LT, LU, LV, MC, MT, NL, PL, PT, RO, SE, SI, SK, TR), OAPI (BF, BJ, CF, CG, CI, CM, GA, GN, GQ, GW, ML, MR, NE, SN, TD, TG).

Published:

— without international search report and to be republished upon receipt of that report

For two-letter codes and other abbreviations, refer to the "Guidance Notes on Codes and Abbreviations" appearing at the beginning of each regular issue of the PCT Gazette.

(54) Title: RNA SILENCING COMPOSITIONS AND METHODS FOR THE TREATMENT OF HUNTINGTON'S DISEASE

(57) Abstract: The present invention relates to the discovery of an effective treatment for a variety of Huntington's disease (HD). The present invention utilizes RNA silencing technology (e.g. RNAi) against single nucleotide polymorphisms (SNPs) in the Huntingtin (htt) gene encoding the dominant, gain-of-function mutant Huntington protein, thereby resulting in an effective treatment for the gain-of-function disease.

WO 2008/005562 A2

RNA SILENCING COMPOSITIONS AND METHODS FOR THE TREATMENT OF HUNTINGTON'S DISEASE

Related Applications

This application claims the benefit of priority to US Provisional Patent Application No. 60/819,704, filed July 7, 2006, entitled "RNA silencing compositions and methods for the treatment of Huntington's Disease".

The content the above identified application is hereby incorporated by reference in it's entirety.

Statement as to Federally Funded Research

The U.S. government may have certain rights in this invention pursuant to Grant No NIH NS38194, awarded by the National Institute of Health (NIH).

Background of the Invention

RNA silencing refers to a group of sequence-specific regulatory mechanisms (e.g. RNA interference (RNAi), transcriptional gene silencing (TGS), post-transcriptional gene silencing (PTGS), quelling, co-suppression, and translational repression) mediated by RNA silencing agents which result in repression or "silencing" of a corresponding protein-coding gene. RNA silencing has been observed in many types of eukaryotes, including humans, and utility of RNA silencing agents as both therapeutics and research tools is the subject of intense interest.

Several types of small (~19-23 nt), noncoding RNAs trigger RNA silencing in eukaryotes, including small interfering RNAs (siRNAs) and microRNAs (miRNAs, also known as small temporal RNAs (stRNAs)). Recent evidence suggests that the two classes of small RNAs are functionally interchangeable, with the choice of RNA silencing mechanism (e.g. RNAi-mediated mRNA cleavage or translational repression) determined largely by the degree of complementarity between the small RNA and its target (Schwarz and Zamore, 2002; Hutvagner and Zamore, 2002; Zeng et al., 2003; Doench et al., 2003). RNA silencing agents with a high degree of complementarity to a corresponding target mRNA have been shown to direct its silencing by the cleavage-based mechanism (Zamore et al., 2000; Elbashir et al., 2001a; Rhoades et al., 2002; Reinhart et al., 2002; Llave et al., 2002a; Llave et al., 2002b; Xie et al., 2003; Kasschau

et al., 2003; Tang et al., 2003; Chen, 2003). RNA silencing agents with a lower degree of complementarity mediate gene silencing by recruiting the RISC complex to the target mRNA, thereby blocking its translation but leaving the mRNA intact (Mourelatos et al., 2002; Hutvagner and Zamore, 2002; Caudy et al., 2002; Martinez et al., 2002; Abrahante et al., 2003; Brennecke et al., 2003; Lin et al., 2003; Xu et al., 2003).

RNA silencing agents have received particular interest as research tools and therapeutic agents for their ability to knock down expression of a particular protein with a high degree of sequence specificity. The sequence specificity of RNA silencing agents is particularly useful for allele-specific silencing dominant, gain-of-function gene mutations. Diseases caused by dominant, gain-of-function gene mutations develop in heterozygotes bearing one mutant and one wild type copy of the gene. Some of the best-known diseases of this class are common neurodegenerative diseases, including Alzheimer's disease, Parkinson's disease and amyotrophic lateral sclerosis (ALS; "Lou Gehrig's disease") (Taylor et al., 2002). In these diseases, the exact pathways whereby the mutant proteins cause cell degeneration are not clear, but the origin of the cellular toxicity is known to be the mutant protein.

One group of inherited gain-of-function disorders are known as the trinucleotide repeat diseases. The common genetic mutation among these diseases is an increase in a series of a particular trinucleotide repeat. To date, the most frequent trinucleotide repeat is CAG, which codes for the amino acid glutamine. At least 9 CAG repeat diseases are known and there are more than 20 varieties of these diseases, including Huntington's disease, Kennedy's disease and many spinocerebellar diseases. These disorders share a neurodegenerative component in the brain and/or spinal cord. Each disease has a specific pattern of neurodegeneration in the brain and most have an autosomal dominant inheritance. The onset of the diseases generally occurs at 30 to 40 years of age, but in Huntington's disease CAG repeats in the huntingtin gene of >60 portend a juvenile onset. Research has shown that the genetic mutation (increase in length of CAG repeats from normal <36 in the huntingtin gene to >36 in disease) is associated with the synthesis of a mutant huntingtin protein, which has >36 polyglutamines (Aronin et al., 1995). It has also been shown that the protein forms cytoplasmic aggregates and nuclear inclusions (Difiglia et al., 1997) and associates with vesicles (Aronin et al., 1999). The precise pathogenic pathways are not known.

In the search for an effective treatment for these diseases, researchers in this field emphasized understanding the pathogenesis of the disease and initially sought to intercede at the level of the presumed aberrant protein interactions. However, there is no approved treatment for Huntington's disease or other trinucleotide repeat diseases. Accordingly, therapeutic RNA silencing agents capable of silencing Huntingtin proteins are of considerable interest.

Summary of the Invention

The present invention is based, at least in part, on the discovery of single nucleotide polymorphism (SNP) sites in the Huntingtin (htt) gene which are preferred target sites for RNA silencing. The htt SNP sites of the invention are relatively prevalent within a sample population. In particular, the htt SNPs of the invention are present within a population at an allelic frequency of at least 30%. Such SNPs sites may be analysed in a patient to determine if they are heterozygous. Each SNP allele of a heterozygous SNP site may then be sequenced in the patient to determine which SNP allele is linked with the expanded CAG repeat region of the HD-associated allele to form a HD-associated haplotype. Such HD-associated htt SNPs are attractive targets for therapeutic RNA silencing agents and circumvent complications associated with directly targeting the expanded CAG repeat region of htt. Over 80 normal genes with CAG repeat regions are known to exist in cells. Thus, RNA silencing agents targeting these CAG repeats cannot be used without risking widespread destruction of normal CAG repeat-containing mRNAs. Likewise, targeting non-allele-specific sites would result in loss of both normal and mutant huntingtin causes neuronal dysfunction.

In one aspect, the invention is directed to a method of treating a subject having or at risk for Huntington's disease, comprising: administering to said subject an effective amount of a RNA silencing agent targeting a heterozygous single nucleotide polymorphism (SNP) within a target mRNA encoding a mutant huntingtin (htt) protein, such that RNA silencing of said mRNA occurs; thereby treating said disease in said subject, wherein the SNP has an allelic frequency of at least 10% (*e.g.*, at least 15%, 20%, 25%, 30%, 35%, 40% or more) in a sample population.

In another aspect, the invention is directed to a method of silencing a target mRNA encoding a mutant huntingtin (htt) protein in a cell, comprising contacting the cell with effective amount of a RNA silencing agent targeting a heterozygous single nucleotide polymorphism (SNP) within the target mRNA, such that RNA silencing of

said mRNA occurs, wherein the SNP has an allelic frequency of at least 10% (*e.g.*, at least 15%, 20%, 25%, 30%, 35%, 40% or more) in a sample population.

In another aspect, the invention is directed to an RNA silencing agent comprising an antisense strand comprising about 16-25 nucleotides homologous to a region of an mRNA encoding a mutant huntingtin (htt) protein, said region comprising a heterozygous single nucleotide polymorphism (SNP) allele having an allelic frequency of at least 10% (*e.g.*, at least 15%, 20%, 25%, 30%, 35%, 40% or more) in a sample population, wherein the RNA silencing agent is capable of directing RNA silencing of said mRNA. In certain embodiments, the heterozygous SNP allele is found at a SNP site selected from the group consisting of RS362331, RS4690077, RS363125, 47bp into Exon 25, RS363075, RS362268, RS362267, RS362307, RS362306, RS362305, RS362304, and RS362303.

In one embodiment, the SNP allele is present at SNP target site RS363125. In a particular embodiment, the SNP allele is a C nucleotide. In another particular embodiment, the SNP allele is a U nucleotide.

In another embodiment, the SNP allele is present at SNP target site RS362331. In a particular embodiment, the SNP allele is an A nucleotide. In another particular embodiment, the SNP allele is a C nucleotide.

In one embodiment, the target mRNA comprises the sequence set forth as SEQ ID NO: 5. In another embodiment, the target mRNA comprises the sequence set forth as SEQ ID NO: 6. In another embodiment, the target mRNA comprises the sequence set forth as SEQ ID NO: 11. In another embodiment, the target mRNA comprises the sequence set forth as SEQ ID NO: 12. In another embodiment, the target mRNA comprises the sequence set forth as SEQ ID NO: 17.

In certain embodiments, the RNA silencing agent is capable of inducing discriminatory RNA silencing.

In one embodiment, the antisense strand of said RNA silencing agent is complementary to the SNP and wherein said RNA silencing agent is capable of substantially silencing the mutant huntingtin protein without substantially silencing the corresponding wild-type huntingtin protein.

In certain preferred embodiments, the RNA silencing agent is an siRNA. In one embodiment, the siRNA comprises (i) a sense strand comprising the sequence set forth as SEQ ID NO: 3; and (ii) an antisense strand comprising the sequence set forth as SEQ ID NO: 4. In another embodiment, the siRNA comprises (i) a sense strand comprising the sequence set forth as SEQ ID NO: 7; and (ii) an antisense strand comprising the sequence set forth as SEQ ID NO: 8. In another embodiment, the siRNA comprises (i) a sense strand comprising the sequence set forth as SEQ ID NO: 9; and (ii) an antisense strand comprising the sequence set forth as SEQ ID NO: 10. In another embodiment, the siRNA comprises (i) a sense strand comprising the sequence set forth as SEQ ID NO: 13; and (ii) an antisense strand comprising the sequence set forth as SEQ ID NO: 14. In another embodiment, the siRNA comprises (i) a sense strand comprising the sequence set forth as SEQ ID NO: 15; and (ii) an antisense strand comprising the sequence set forth as SEQ ID NO: 16. In another embodiment, the siRNA comprises (i) a sense strand comprising the sequence set forth as SEQ ID NO: 18; and (ii) an antisense strand comprising the sequence set forth as SEQ ID NO: 19. In another embodiment, at least one nucleotide of the siRNA is modified with a nucleotide analog or backbone modification (e.g., a phosphorothioate or Locked Nucleic Acid (LNA) modification) which confers, for example, enhanced nuclease resistance.

In other embodiments, the siRNA comprises a lipophilic moiety. In a particular preferred embodiment, the lipophilic moiety is a cholesterol moiety.

In other embodiments, the siRNA is an asymmetric siRNA.

In certain embodiments, the subject is identified as having said SNP by (i) providing DNA from the subject; and (ii) sequencing the huntingtin gene, or portion thereof, using said DNA. In other embodiments, the sample population is of Western European origin.

Other features and advantages of the invention will be apparent from the following detailed description and claims.

Brief Description of the Drawings

Figure 1: Frequency of SNP heterozygosity at SNP sites located in the human Huntingtin gene. The identity of the SNP at each SNP site in the target gene and target mRNA are also indicated.

Figure 2a-b: *In vitro* RNAi reactions programmed with siRNA targeting a first SNP allele (C) in the heterozygous SNP site RS363331 within the human huntingtin gene. (a) Sequence of the siRNA (SEQ ID NO: 3; sense strand; SEQ ID NO: 4, guide strand), which is fully complementary to the target *hht* mRNA containing the "C" SNP allele (SEQ ID NO: 5) but which forms a G:U mismatch at position 10 (P10) with the non-target mRNA encoded by the corresponding "U" SNP (SEQ ID NO: 6). (b) Discriminatory RNA silencing (expressed as units of luciferase reporter gene activity relative to GFP) by the siRNA in (a) for the targeted "U" SNP allele ("match") versus the non-targeted "C" SNP allele ("mismatch").

Figure 3a-b: *In vitro* RNAi reactions programmed with siRNA targeting a second SNP allele (T) at heterozygous SNP site RS363331 within the human huntingtin gene. (a) Sequence of the siRNA (SEQ ID NO: 7; sense strand; SEQ ID NO: 8, guide strand) which is fully complementary to the target *hht* mRNA containing the "U" SNP allele (SEQ ID NO: 6) but which forms a A:C mismatch at position 10 (P10) with the non-target mRNA encoded by the corresponding "C" SNP (SEQ ID NO: 5). (b) Discriminatory RNA silencing (expressed as units of luciferase reporter gene activity relative to GFP) by the siRNA in (a) for the targeted "U" SNP allele ("match") versus the non-targeted "C" SNP allele ("mismatch").

Figure 4a-b: *In vitro* RNAi reactions programmed with siRNA targeting a first SNP allele (A) in the heterozygous SNP site RS363125 within the human huntingtin gene. (a) Sequence of the siRNA (SEQ ID NO: 9; sense strand; SEQ ID NO: 10, guide strand), which is fully complementary to the target *hht* mRNA containing the "A" SNP allele (SEQ ID NO: 11) but which forms a U:C mismatch at position 10 (P10) with the non-target mRNA encoded by the corresponding "C" SNP (SEQ ID NO: 12). (b) Discriminatory RNA silencing (expressed as units of luciferase reporter gene activity relative to GFP) by the siRNA in (a) for the targeted "A" SNP allele ("match") versus the non-targeted "C" SNP allele ("mismatch").

Figure 5a-b: *In vitro* RNAi reactions programmed with siRNA targeting the second SNP allele ("C") at the heterozygous SNP site RS363125 within the human

huntingtin gene. (a) Sequence of the siRNA (SEQ ID NO: 13; sense strand; SEQ ID NO: 14, guide strand) which is fully complementary to the target *htt* mRNA containing the "C" SNP allele (SEQ ID NO: 12) but which forms a G:A mismatch at position 10 (P10) with the non-target mRNA encoded by the corresponding "C" SNP (SEQ ID NO: 11). (b) Discriminatory RNA silencing (expressed as units of luciferase reporter gene activity relative to GFP) by the siRNA in (a) for the targeted "C" SNP allele ("match") versus the non-targeted "A" SNP allele ("mismatch").

Figure 6a-d: *In vitro* RNAi reactions performed in HEK cells homozygous for the C polymorphism in the 3' UTR of the human huntingtin (*htt*) gene. (a) Sequence of a matched siRNA (SEQ ID NO: 15, sense strand; SEQ ID NO: 16, guide strand) having a guide strand that is perfectly complementary to the target site in the homozygous target allele (SEQ ID NO: 17). (b) Sequence of a mismatched siRNA (SEQ ID NO: 18, sense strand; SEQ ID NO: 19, guide strand) having a guide strand which forms a U:C mismatch at position 10 (P10) with the homozygous target allele. (c) Relative change in *htt* target mRNA levels in HEK cells transfected with 20nM of siRNAs depicted in (a) and (b) versus mock transfection as measured by quantitative RT-PCR. (d) Relative change in Htt protein levels in HEK cells transfected with 20nM of siRNAs depicted in (a) and (b) versus mock transfection as measured by Western blot.

Figure 7: Relative change in *htt* target mRNA levels in HEK cells transfected with 5, 10, or 20nM of the matched and mismatched siRNAs depicted in Figure 6(a) and 6(b) and an unrelated GFP siRNA as measured by quantitative RT-PCR.

Detailed Description of the Invention

The present invention relates to methods and reagents for treating a variety of gain-of-function diseases. In one aspect, the invention relates to methods and reagents for treating a variety of diseases characterized by a mutation in one allele or copy of a gene, the mutation encoding a protein which is sufficient to contribute to or cause the disease. Preferably, the methods and reagents are used to treat Huntington's Disease.

The present invention utilizes RNA silencing technology (e.g. RNAi) against single nucleotide polymorphisms (SNPs) located within the *htt* gene encoding the mutant Huntington protein. RNA silencing destroys the corresponding mutant mRNA with single nucleotide specificity and selectivity. RNA silencing agents of the present invention are targeted to polymorphic regions of the mutant *htt* gene, resulting in

cleavage or translational repression of mutant *htt* mRNA. These RNA silencing agents, through a series of protein-nucleotide interactions, function to cleave or translationally repress the mutant *htt* mRNAs.

Definitions

So that the invention may be more readily understood, certain terms are first defined.

As used herein, the term "RNA silencing" refers to a group of sequence-specific regulatory mechanisms (e.g. RNA interference (RNAi), transcriptional gene silencing (TGS), post-transcriptional gene silencing (PTGS), quelling, co-suppression, and translational repression) mediated by RNA molecules which result in the inhibition or "silencing" of the expression of a corresponding protein-coding gene. RNA silencing has been observed in many types of organisms, including plants, animals, and fungi.

The term "discriminatory RNA silencing" refers to the ability of an RNA molecule to substantially inhibit the expression of a "first" or "target" polynucleotide sequence while not substantially inhibiting the expression of a "second" or "non-target" polynucleotide sequence", *e.g.*, when both polynucleotide sequences are present in the same cell. In certain embodiments, the target polynucleotide sequence corresponds to a target gene, while the non-target polynucleotide sequence corresponds to a non-target gene. In other embodiments, the target polynucleotide sequence corresponds to a target allele, while the non-target polynucleotide sequence corresponds to a non-target allele. In certain embodiments, the target polynucleotide sequence is the DNA sequence encoding the regulatory region (*e.g.* promoter or enhancer elements) of a target gene. In other embodiments, the target polynucleotide sequence is a target mRNA encoded by a target gene.

As used herein, the term "target gene" is a gene whose expression is to be substantially inhibited or "silenced." This silencing can be achieved by RNA silencing, *e.g.* by cleaving the mRNA of the target gene or translational repression of the target gene. The term "non-target gene" is a gene whose expression is not to be substantially silenced. In one embodiment, the polynucleotide sequences of the target and non-target gene (*e.g.* mRNA encoded by the target and non-target genes) can differ by one or more

nucleotides. In another embodiment, the target and non-target genes can differ by one or more polymorphisms (*e.g.*, Single Nucleotide Polymorphisms or SNPs). In another embodiment, the target and non-target genes can share less than 100% sequence identity. In another embodiment, the non-target gene may be a homolog (*e.g.* an ortholog or paralog) of the target gene.

A “target allele” is an allele (*e.g.*, a SNP allele) whose expression is to be selectively inhibited or “silenced.” This silencing can be achieved by RNA silencing, *e.g.* by cleaving the mRNA of the target gene or target allele by a siRNA. The term “non-target allele” is a allele whose expression is not to be substantially silenced. In certain embodiments, the target and non-target alleles can correspond to the same target gene. In other embodiments, the target allele corresponds to, or is associated with, a target gene, and the non-target allele corresponds to, or is associated with, a non-target gene. In one embodiment, the polynucleotide sequences of the target and non-target alleles can differ by one or more nucleotides. In another embodiment, the target and non-target alleles can differ by one or more allelic polymorphisms (*e.g.*, one or more SNPs). In another embodiment, the target and non-target alleles can share less than 100% sequence identity.

The term “polymorphism” as used herein, refers to a variation (*e.g.*, one or more deletions, insertions, or substitutions) in a gene sequence that is identified or detected when the same gene sequence from different sources or subjects (but from the same organism) are compared. For example, a polymorphism can be identified when the same gene sequence from different subjects are compared. Identification of such polymorphisms is routine in the art, the methodologies being similar to those used to detect, for example, breast cancer point mutations. Identification can be made, for example, from DNA extracted from a subject’s lymphocytes, followed by amplification of polymorphic regions using specific primers to said polymorphic region. Alternatively, the polymorphism can be identified when two alleles of the same gene are compared. In particular embodiments, the polymorphism is a single nucleotide polymorphism (SNP).

A variation in sequence between two alleles of the same gene within an organism is referred to herein as an “allelic polymorphism”. In certain embodiments, the allelic polymorphism corresponds to a SNP allele. For example, the allelic polymorphism may comprise a single nucleotide variation between the two alleles of a SNP. The

polymorphism can be at a nucleotide within a coding region but, due to the degeneracy of the genetic code, no change in amino acid sequence is encoded. Alternatively, polymorphic sequences can encode a different amino acid at a particular position, but the change in the amino acid does not affect protein function. Polymorphic regions can also be found in non-encoding regions of the gene. In preferred embodiments, the polymorphism is found in a coding region of the gene or in an untranslated region (*e.g.*, a 5' UTR or 3' UTR) of the gene.

As used herein, the term "allelic frequency" is a measure (*e.g.*, proportion or percentage) of the relative frequency of an allele (*e.g.*, a SNP allele) at a single locus in a population of individuals. For example, where a population of individuals carry n loci of a particular chromosomal locus (and the gene occupying the locus) in each of their somatic cells, then the allelic frequency of an allele is the fraction or percentage of loci that the allele occupies within the population. In particular embodiments, the allelic frequency of an allele (*e.g.* a SNP allele) is at least 10% (*e.g.*, at least 15%, 20%, 25%, 30%, 35%, 40% or more) in a sample population.

As used herein, the term "sample population" refers to a population of individuals comprising a statistically significant number of individuals. For example, the sample population may comprise 50, 75, 100, 200, 500, 1000 or more individuals. In particular embodiments, the sample population may comprise individuals which share at least one common disease phenotype (*e.g.*, a gain-of-function disorder) or mutation (*e.g.*, a gain-of-function mutation).

As used herein, the term "heterozygosity" refers to the fraction of individuals within a population that are heterozygous (*e.g.*, contain two or more different alleles) at a particular locus (*e.g.*, at a SNP). Heterozygosity may be calculated for a sample population using methods that are well known to those skilled in the art.

The term "gain-of-function mutation" as used herein, refers to any mutation in a gene in which the protein encoded by said gene (*i.e.*, the mutant protein) acquires a function not normally associated with the protein (*i.e.*, the wild type protein) causes or contributes to a disease or disorder. The gain-of-function mutation can be a deletion, addition, or substitution of a nucleotide or nucleotides in the gene which gives rise to the change in the function of the encoded protein. In one embodiment, the gain-of-function mutation changes the function of the mutant protein or causes interactions with other proteins. In another embodiment, the gain-of-function mutation causes a decrease in or

removal of normal wild-type protein, for example, by interaction of the altered, mutant protein with said normal, wild-type protein.

As used herein, the term “gain-of-function disorder”, refers to a disorder characterized by a gain-of-function mutation. In one embodiment, the gain-of-function disorder is a neurodegenerative disease caused by a gain-of-function mutation, e.g., polyglutamine disorders and/or trinucleotide repeat diseases, for example, Huntington’s disease. In another embodiment, the gain-of-function disorder is caused by a gain-of-function in an oncogene, the mutated gene product being a gain-of-function mutant, e.g., cancers caused by a mutation in the *ret* oncogene (e.g., *ret-1*), for example, endocrine tumors, medullary thyroid tumors, parathyroid hormone tumors, multiple endocrine neoplasia type2, and the like. Additional exemplary gain-of-function disorders include Alzheimer’s, human immunodeficiency disorder (HIV), and slow channel congenital myasthenic syndrome (SCCMS).

The term “trinucleotide repeat diseases” as used herein, refers to any disease or disorder characterized by an expanded trinucleotide repeat region located within a gene, the expanded trinucleotide repeat region being causative of the disease or disorder. Examples of trinucleotide repeat diseases include, but are not limited to spino-cerebellar ataxia type 12 spino-cerebellar ataxia type 8, fragile X syndrome, fragile XE Mental Retardation, Friedreich’s ataxia and myotonic dystrophy. Preferred trinucleotide repeat diseases for treatment according to the present invention are those characterized or caused by an expanded trinucleotide repeat region at the 5’ end of the coding region of a gene, the gene encoding a mutant protein which causes or is causative of the disease or disorder. Certain trinucleotide diseases, for example, fragile X syndrome, where the mutation is not associated with a coding region may not be suitable for treatment according to the methodologies of the present invention, as there is no suitable mRNA to be targeted by RNAi. By contrast, disease such as Friedreich’s ataxia may be suitable for treatment according to the methodologies of the invention because, although the causative mutation is not within a coding region (*i.e.*, lies within an intron), the mutation may be within, for example, an mRNA precursor (e.g., a pre-spliced mRNA precursor).

The term “polyglutamine disorder” as used herein, refers to any disease or disorder characterized by an expanded of a (CAG)_n repeats at the 5’ end of the coding region (thus encoding an expanded polyglutamine region in the encoded protein). In one embodiment, polyglutamine disorders are characterized by a progressive

degeneration of nerve cells. Examples of polyglutamine disorders include but are not limited to: Huntington's disease, spino-cerebellar ataxia type 1, spino-cerebellar ataxia type 2, spino-cerebellar ataxia type 3 (also known as Machado-Joseph disease), and spino-cerebellar ataxia type 6, spino-cerebellar ataxia type 7 and dentato-rubral-pallidolusian atrophy.

The term "polyglutamine domain," as used herein, refers to a segment or domain of a protein that consists of a consecutive glutamine residues linked to peptide bonds. In one embodiment the consecutive region includes at least 5 glutamine residues.

The term "expanded polyglutamine domain" or "expanded polyglutamine segment", as used herein, refers to a segment or domain of a protein that includes at least 35 consecutive glutamine residues linked by peptide bonds. Such expanded segments are found in subjects afflicted with a polyglutamine disorder, as described herein, whether or not the subject has shown to manifest symptoms.

The term "trinucleotide repeat" or "trinucleotide repeat region" as used herein, refers to a segment of a nucleic acid sequence (e.g.,) that consists of consecutive repeats of a particular trinucleotide sequence. In one embodiment, the trinucleotide repeat includes at least 5 consecutive trinucleotide sequences. Exemplary trinucleotide sequences include, but are not limited to, CAG, CGG, GCC, GAA, CTG, and/or CGG.

As used herein, the term "RNA silencing agent" refers to an RNA which is capable of inhibiting or "silencing" the expression of a target gene. In certain embodiments, the RNA silencing agent is capable of preventing complete processing (e.g., the full translation and/or expression) of a mRNA molecule through a post-transcriptional silencing mechanism. RNA silencing agents include small (<50 b.p.), noncoding RNA molecules, for example RNA duplexes comprising paired strands, as well as precursor RNAs from which such small non-coding RNAs can be generated. Exemplary RNA silencing agents include siRNAs, miRNAs, siRNA-like duplexes, and dual-function oligonucleotides as well as precursors thereof. In one embodiment, the RNA silencing agent is capable of inducing RNA interference. In another embodiment, the RNA silencing agent is capable of mediating translational repression.

The term "nucleoside" refers to a molecule having a purine or pyrimidine base covalently linked to a ribose or deoxyribose sugar. Exemplary nucleosides include adenosine, guanosine, cytidine, uridine and thymidine. Additional exemplary nucleosides include inosine, 1-methyl inosine, pseudouridine, 5,6-dihydrouridine,

ribothymidine, ^2N -methylguanosine and $^{2,2}\text{N}$,N-dimethylguanosine (also referred to as "rare" nucleosides). The term "nucleotide" refers to a nucleoside having one or more phosphate groups joined in ester linkages to the sugar moiety. Exemplary nucleotides include nucleoside monophosphates, diphosphates and triphosphates. The terms "polynucleotide" and "nucleic acid molecule" are used interchangeably herein and refer to a polymer of nucleotides joined together by a phosphodiester linkage between 5' and 3' carbon atoms.

The term "RNA" or "RNA molecule" or "ribonucleic acid molecule" refers to a polymer of ribonucleotides (*e.g.*, 2, 3, 4, 5, 10, 15, 20, 25, 30, or more ribonucleotides). The term "DNA" or "DNA molecule" or "deoxyribonucleic acid molecule" refers to a polymer of deoxyribonucleotides. DNA and RNA can be synthesized naturally (*e.g.*, by DNA replication or transcription of DNA, respectively). RNA can be post-transcriptionally modified. DNA and RNA can also be chemically synthesized. DNA and RNA can be single-stranded (*i.e.*, ssRNA and ssDNA, respectively) or multi-stranded (*e.g.*, double stranded, *i.e.*, dsRNA and dsDNA, respectively). "mRNA" or "messenger RNA" is single-stranded RNA that specifies the amino acid sequence of one or more polypeptide chains. This information is translated during protein synthesis when ribosomes bind to the mRNA.

As used herein, the term "rare nucleotide" refers to a naturally occurring nucleotide that occurs infrequently, including naturally occurring deoxyribonucleotides or ribonucleotides that occur infrequently, *e.g.*, a naturally occurring ribonucleotide that is not guanosine, adenosine, cytosine, or uridine. Examples of rare nucleotides include, but are not limited to, inosine, 1-methyl inosine, pseudouridine, 5,6-dihydrouridine, ribothymidine, ^2N -methylguanosine and $^{2,2}\text{N}$,N-dimethylguanosine.

The term "engineered," as in an engineered RNA precursor, or an engineered nucleic acid molecule, indicates that the precursor or molecule is not found in nature, in that all or a portion of the nucleic acid sequence of the precursor or molecule is created or selected by man. Once created or selected, the sequence can be replicated, translated, transcribed, or otherwise processed by mechanisms within a cell. Thus, an RNA precursor produced within a cell from a transgene that includes an engineered nucleic acid molecule is an engineered RNA precursor.

As used herein, the term "small interfering RNA" ("siRNA") (also referred to in the art as "short interfering RNAs") refers to an RNA (or RNA analog) comprising

between about 10-50 nucleotides (or nucleotide analogs) which is capable of directing or mediating RNA interference. Preferably, a siRNA comprises between about 15-30 nucleotides or nucleotide analogs, more preferably between about 16-25 nucleotides (or nucleotide analogs), even more preferably between about 18-23 nucleotides (or nucleotide analogs), and even more preferably between about 19-22 nucleotides (or nucleotide analogs) (e.g., 19, 20, 21 or 22 nucleotides or nucleotide analogs). The term "short" siRNA refers to a siRNA comprising ~21 nucleotides (or nucleotide analogs), for example, 19, 20, 21 or 22 nucleotides. The term "long" siRNA refers to a siRNA comprising ~24-25 nucleotides, for example, 23, 24, 25 or 26 nucleotides. Short siRNAs may, in some instances, include fewer than 19 nucleotides, e.g., 16, 17 or 18 nucleotides, provided that the shorter siRNA retains the ability to mediate RNAi. Likewise, long siRNAs may, in some instances, include more than 26 nucleotides, provided that the longer siRNA retains the ability to mediate RNAi absent further processing, e.g., enzymatic processing, to a short siRNA.

As used herein, the term "microRNA" ("miRNA"), also referred to in the art as "small temporal RNAs" ("stRNAs"), refers to a small (10-50 nucleotide) RNA which are genetically encoded (e.g. by viral, mammalian, or plant genomes) and are capable of directing or mediating RNA silencing. An "miRNA disorder" shall refer to a disease or disorder characterized by an aberrant expression or activity of an miRNA.

As used herein, the term "dual functional oligonucleotide" refers to a RNA silencing agent having the formula $T-L-\mu$, wherein T is an mRNA targeting moiety, L is a linking moiety, and μ is a miRNA recruiting moiety. As used herein, the terms "mRNA targeting moiety", "targeting moiety", "mRNA targeting portion" or "targeting portion" refer to a domain, portion or region of the dual functional oligonucleotide having sufficient size and sufficient complementarity to a portion or region of an mRNA chosen or targeted for silencing (*i.e.*, the moiety has a sequence sufficient to capture the target mRNA). As used herein, the term "linking moiety" or "linking portion" refers to a domain, portion or region of the RNA-silencing agent which covalently joins or links the mRNA

As used herein, the term "antisense strand" of an RNA silencing agent, *e.g.* an siRNA or RNA silencing agent, refers to a strand that is substantially complementary to a section of about 10-50 nucleotides, *e.g.*, about 15-30, 16-25, 18-23 or 19-22 nucleotides of the mRNA of the gene targeted for silencing. The antisense strand or first

strand has sequence sufficiently complementary to the desired target mRNA sequence to direct target-specific silencing, e.g., complementarity sufficient to trigger the destruction of the desired target mRNA by the RNAi machinery or process (RNAi interference) or complementarity sufficient to trigger translational repression of the desired target mRNA.

The term "sense strand" or "second strand" of an RNA silencing agent, *e.g.* an siRNA or RNA silencing agent, refers to a strand that is complementary to the antisense strand or first strand. Antisense and sense strands can also be referred to as first or second strands, the first or second strand having complementarity to the target sequence and the respective second or first strand having complementarity to said first or second strand. miRNA duplex intermediates or siRNA-like duplexes include a miRNA strand having sufficient complementarity to a section of about 10-50 nucleotides of the mRNA of the gene targeted for silencing and a miRNA* strand having sufficient complementarity to form a duplex with the miRNA strand.

As used herein, the term "guide strand" refers to a strand of an RNA silencing agent, *e.g.*, an antisense strand of an siRNA duplex or siRNA sequence, that enters into the RISC complex and directs cleavage of the target mRNA.

The term "nucleotide analog" or "altered nucleotide" or "modified nucleotide" refers to a non-standard nucleotide, including non-naturally occurring ribonucleotides or deoxyribonucleotides. Preferred nucleotide analogs are modified at any position so as to alter certain chemical properties of the nucleotide yet retain the ability of the nucleotide analog to perform its intended function. Examples of positions of the nucleotide which may be derivitized include the 5 position, *e.g.*, 5-(2-amino)propyl uridine, 5-bromo uridine, 5-propyne uridine, 5-propenyl uridine, etc.; the 6 position, *e.g.*, 6-(2-amino)propyl uridine; the 8-position for adenosine and/or guanosines, *e.g.*, 8-bromo guanosine, 8-chloro guanosine, 8-fluoroguanosine, etc. Nucleotide analogs also include deaza nucleotides, *e.g.*, 7-deaza-adenosine; O- and N-modified (*e.g.*, alkylated, *e.g.*, N6-methyl adenosine, or as otherwise known in the art) nucleotides; and other heterocyclically modified nucleotide analogs such as those described in Herdewijn, *Antisense Nucleic Acid Drug Dev.*, 2000 Aug. 10(4):297-310.

Nucleotide analogs may also comprise modifications to the sugar portion of the nucleotides. For example the 2' OH-group may be replaced by a group selected from H,

OR, R, F, Cl, Br, I, SH, SR, NH₂, NHR, NR₂, COOR, or OR, wherein R is substituted or unsubstituted C₁–C₆ alkyl, alkenyl, alkynyl, aryl, etc. Other possible modifications include those described in U.S. Patent Nos. 5,858,988, and 6,291,438.

The phosphate group of the nucleotide may also be modified, e.g., by substituting one or more of the oxygens of the phosphate group with sulfur (e.g., phosphorothioates), or by making other substitutions which allow the nucleotide to perform its intended function such as described in, for example, Eckstein, *Antisense Nucleic Acid Drug Dev.* 2000 Apr. 10(2):117-21, Rusckowski et al. *Antisense Nucleic Acid Drug Dev.* 2000 Oct. 10(5):333-45, Stein, *Antisense Nucleic Acid Drug Dev.* 2001 Oct. 11(5): 317-25, Vorobjev et al. *Antisense Nucleic Acid Drug Dev.* 2001 Apr. 11(2):77-85, and U.S. Patent No. 5,684,143. Certain of the above-referenced modifications (e.g., phosphate group modifications) preferably decrease the rate of hydrolysis of, for example, polynucleotides comprising said analogs in vivo or in vitro.

As used herein, the term “asymmetry”, as in the asymmetry of the duplex region of an RNA silencing agent (e.g. the stem of an shRNA), refers to an inequality of bond strength or base pairing strength between the termini of the RNA silencing agent (e.g., between terminal nucleotides on a first strand or stem portion and terminal nucleotides on an opposing second strand or stem portion), such that the 5' end of one strand of the duplex is more frequently in a transient unpaired, e.g. single-stranded, state than the 5' end of the complementary strand. This structural difference determines that one strand of the duplex is preferentially incorporated into a RISC complex. The strand whose 5' end is less tightly paired to the complementary strand will preferentially be incorporated into RISC and mediate RNAi.

As used herein, the term “bond strength” or “base pair strength” refers to the strength of the interaction between pairs of nucleotides (or nucleotide analogs) on opposing strands of an oligonucleotide duplex (e.g., an siRNA duplex), due primarily to H-bonding, Van der Waals interactions, and the like between said nucleotides (or nucleotide analogs).

As used herein, the “5' end”, as in the 5' end of an antisense strand, refers to the 5' terminal nucleotides, e.g., between one and about 5 nucleotides at the 5' terminus of the antisense strand. As used herein, the “3' end”, as in the 3' end of a sense strand, refers to the region, e.g., a region of between one and about 5 nucleotides, that is complementary to the nucleotides of the 5' end of the complementary antisense strand.

As used herein the term “destabilizing nucleotide” refers to a first nucleotide or nucleotide analog capable of forming a base pair with second nucleotide or nucleotide analog such that the base pair is of lower bond strength than a conventional base pair (ie. Watson-Crick base pair). In certain embodiments, the destabilizing nucleotide is capable of forming a mismatch base pair with the second nucleotide. In other embodiments, the destabilizing nucleotide is capable of forming a wobble base pair with the second nucleotide. In yet other embodiments, the destabilizing nucleotide is capable of forming an ambiguous base pair with the second nucleotide.

As used herein, the term “base pair” refers to the interaction between pairs of nucleotides (or nucleotide analogs) on opposing strands of an oligonucleotide duplex (e.g., a duplex formed by a strand of a RNA silencing agent and a target mRNA sequence), due primarily to H-bonding, Van der Waals interactions, and the like between said nucleotides (or nucleotide analogs). As used herein, the term “bond strength” or “base pair strength” refers to the strength of the base pair.

As used herein, the term “mismatched base pair” refers to a base pair consisting of noncomplementary or non-Watson-Crick base pairs, for example, not normal complementary G:C, A:T or A:U base pairs. As used herein the term “ambiguous base pair” (also known as a non-discriminatory base pair) refers to a base pair formed by a universal nucleotide.

As used herein, term “universal nucleotide” (also known as a “neutral nucleotide”) include those nucleotides (e.g. certain destabilizing nucleotides) having a base (a “universal base” or “neutral base”) that does not significantly discriminate between bases on a complementary polynucleotide when forming a base pair. Universal nucleotides are predominantly hydrophobic molecules that can pack efficiently into antiparallel duplex nucleic acids (e.g. double-stranded DNA or RNA) due to stacking interactions. The base portion of universal nucleotides typically comprise a nitrogen-containing aromatic heterocyclic moiety.

As used herein, the terms “sufficient complementarity” or “sufficient degree of complementarity” mean that the RNA silencing agent has a sequence (e.g. in the antisense strand, mRNA targeting moiety or miRNA recruiting moiety) which is sufficient to bind the desired target RNA, respectively, and to trigger the RNA silencing of the target mRNA.

The term “oligonucleotide” refers to a short polymer of nucleotides and/or nucleotide analogs. The term “RNA analog” refers to a polynucleotide (e.g., a chemically synthesized polynucleotide) having at least one altered or modified nucleotide as compared to a corresponding unaltered or unmodified RNA but retaining the same or similar nature or function as the corresponding unaltered or unmodified RNA. As discussed above, the oligonucleotides may be linked with linkages which result in a lower rate of hydrolysis of the RNA analog as compared to an RNA molecule with phosphodiester linkages. For example, the nucleotides of the analog may comprise methylenediol, ethylene diol, oxymethylthio, oxyethylthio, oxycarbonyloxy, phosphorodiamidate, phosphoroamidate, and/or phosphorothioate linkages. Preferred RNA analogues include sugar- and/or backbone-modified ribonucleotides and/or deoxyribonucleotides. Such alterations or modifications can further include addition of non-nucleotide material, such as to the end(s) of the RNA or internally (at one or more nucleotides of the RNA). An RNA analog need only be sufficiently similar to natural RNA that it has the ability to mediate (mediates) RNA interference.

As used herein, the term “RNA interference” (“RNAi”) refers to a selective intracellular degradation of RNA. RNAi occurs in cells naturally to remove foreign RNAs (e.g., viral RNAs). Natural RNAi proceeds via fragments cleaved from free dsRNA which direct the degradative mechanism to other similar RNA sequences. Alternatively, RNAi can be initiated by the hand of man, for example, to silence the expression of target genes.

As used herein, the term “translational repression” refers to a selective inhibition of mRNA translation. Natural translational repression proceeds *via* miRNAs cleaved from shRNA precursors. Both RNAi and translational repression are mediated by RISC. Both RNAi and translational repression occur naturally or can be initiated by the hand of man, for example, to silence the expression of target genes.

An RNA silencing agent having a strand which is “sequence sufficiently complementary to a target mRNA sequence to direct target-specific RNA interference (RNAi)” means that the strand has a sequence sufficient to trigger the destruction of the target mRNA by the RNAi machinery or process.

As used herein, the term “isolated RNA” (e.g., “isolated siRNA” or “isolated siRNA precursor”) refers to RNA molecules which are substantially free of other cellular material, or culture medium when produced by recombinant techniques, or

substantially free of chemical precursors or other chemicals when chemically synthesized.

The term “in vitro” has its art recognized meaning, e.g., involving purified reagents or extracts, e.g., cell extracts. The term “in vivo” also has its art recognized meaning, e.g., involving living cells, e.g., immortalized cells, primary cells, cell lines, and/or cells in an organism.

As used herein, the term “transgene” refers to any nucleic acid molecule, which is inserted by artifice into a cell, and becomes part of the genome of the organism that develops from the cell. Such a transgene may include a gene that is partly or entirely heterologous (*i.e.*, foreign) to the transgenic organism, or may represent a gene homologous to an endogenous gene of the organism. The term “transgene” also means a nucleic acid molecule that includes one or more selected nucleic acid sequences, e.g., DNAs, that encode one or more engineered RNA precursors, to be expressed in a transgenic organism, e.g., animal, which is partly or entirely heterologous, *i.e.*, foreign, to the transgenic animal, or homologous to an endogenous gene of the transgenic animal, but which is designed to be inserted into the animal's genome at a location which differs from that of the natural gene. A transgene includes one or more promoters and any other DNA, such as introns, necessary for expression of the selected nucleic acid sequence, all operably linked to the selected sequence, and may include an enhancer sequence.

The phrase “examining the function of a gene in a cell or organism” refers to examining or studying the expression, activity, function or phenotype arising therefrom.

Various methodologies of the instant invention include step that involves comparing a value, level, feature, characteristic, property, etc. to a “suitable control”, referred to interchangeably herein as an “appropriate control”. A “suitable control” or “appropriate control” is any control or standard familiar to one of ordinary skill in the art useful for comparison purposes. In one embodiment, a “suitable control” or “appropriate control” is a value, level, feature, characteristic, property, etc. determined prior to performing an RNAi methodology, as described herein. For example, a transcription rate, mRNA level, translation rate, protein level, biological activity, cellular characteristic or property, genotype, phenotype, etc. can be determined prior to introducing an RNA silencing agent of the invention into a cell or organism. In another embodiment, a “suitable control” or “appropriate control” is a value, level, feature,

characteristic, property, etc. determined in a cell or organism, e.g., a control or normal cell or organism, exhibiting, for example, normal traits. In yet another embodiment, a "suitable control" or "appropriate control" is a predefined value, level, feature, characteristic, property, etc.

Unless otherwise defined, all technical and scientific terms used herein have the same meaning as commonly understood by one of ordinary skill in the art to which this invention belongs. Although methods and materials similar or equivalent to those described herein can be used in the practice or testing of the present invention, suitable methods and materials are described below. All publications, patent applications, patents, and other references mentioned herein are incorporated by reference in their entirety. In case of conflict, the present specification, including definitions, will control. In addition, the materials, methods, and examples are illustrative only and not intended to be limiting.

Various aspects of the invention are described in further detail in the following subsections.

I. Huntington Disease

The RNA silencing agents of the invention are designed to target polymorphisms (e.g. single nucleotide polymorphisms) in the mutant human huntingtin protein (*htt*) for the treatment of Huntington's disease.

Huntington's disease, inherited as an autosomal dominant disease, causes impaired cognition and motor disease. Patients can live more than a decade with severe debilitation, before premature death from starvation or infection. The disease begins in the fourth or fifth decade for most cases, but a subset of patients manifest disease in teenage years. The genetic mutation for Huntington's disease is a lengthened CAG repeat in the huntingtin gene. CAG repeat varies in number from 8 to 35 in normal individuals (Kremer et al., 1994). The genetic mutation e.g., an increase in length of the CAG repeats from normal less than 36 in the huntingtin gene to greater than 36 in the disease is associated with the synthesis of a mutant huntingtin protein, which has greater than 36 polyglutamates (Aronin et al., 1995). In general, individuals with 36 or more CAG repeats will get Huntington's disease. Prototypic for as many as twenty other diseases with a lengthened CAG as the underlying mutation, Huntington's disease still has no effective therapy. A variety of interventions -- such as interruption of apoptotic

pathways, addition of reagents to boost mitochondrial efficiency, and blockade of NMDA receptors -- have shown promise in cell cultures and mouse model of Huntington's disease. However, at best these approaches reveal a short prolongation of cell or animal survival.

Huntington's disease complies with the central dogma of genetics: a mutant gene serves as a template for production of a mutant mRNA; the mutant mRNA then directs synthesis of a mutant protein (Aronin et al., 1995; DiFiglia et al., 1997). Mutant huntingtin (protein) probably accumulates in selective neurons in the striatum and cortex, disrupts as yet determined cellular activities, and causes neuronal dysfunction and death (Aronin et al., 1999; Laforet et al., 2001). Because a single copy of a mutant gene suffices to cause Huntington's disease, the most parsimonious treatment would render the mutant gene ineffective. Theoretical approaches might include stopping gene transcription of mutant huntingtin, destroying mutant mRNA, and blocking translation. Each has the same outcome -- loss of mutant huntingtin.

The disease gene linked to Huntington's disease is termed Huntington or (htt). The huntingtin locus is large, spanning 180 kb and consisting of 67 exons. The huntingtin gene is widely expressed and is required for normal development. It is expressed as 2 alternatively polyadenylated forms displaying different relative abundance in various fetal and adult tissues. The larger transcript is approximately 13.7 kb and is expressed predominantly in adult and fetal brain whereas the smaller transcript of approximately 10.3 kb is more widely expressed. The two transcripts differ with respect to their 3' untranslated regions (Lin et al., 1993). Both messages are predicted to encode a 348 kilodalton protein containing 3144 amino acids. The genetic defect leading to Huntington's disease is believed to confer a new property on the mRNA or alter the function of the protein. The amino acid sequence of the human huntingtin protein is set as SEQ ID NO: 2.

A consensus nucleotide sequence of the human huntingtin gene (cDNA) is set forth as SEQ ID NO: 1. The coding region consists of nucleotides 316 to 9750 of SEQ ID NO:1. The two alternative polyadenylation signals are found at nucleotides 10326 to 10331 and nucleotides 13644 to 13649, respectively. The corresponding two polyadenylation sites are found at nucleotides 10348 and 13672, respectively. The first polyadenylation signal/site is that of the 10.3 kb transcript. The second polyadenylation signal/site is that of the 13.7 kb transcript, the predominant transcript in brain.

i. Huntington SNPs

Exemplary single nucleotide polymorphisms in the huntingtin gene sequence can be found at positions 2886, 4034, 6912, 7222, and 7246 of the human htt gene. In preferred embodiments, RNA silencing agents of the invention are capable of targeting one of the SNP sites listed in Figure 1. Genomic sequence for each SNP site can be found in, for example, the publically available "SNP Entrez" database maintained by the NCBI. Additional single nucleotide polymorphisms in the huntingtin gene sequence are set forth in Table 1 below.

Table 1. Exemplary SNPs in the Huntingtin gene.

	consensus	polymorphism	db_xref
complement 103	G	A P6	dbSNP:396875
complement 432	T	C P7	dbSNP:473915
complement 474	C	A P8	dbSNP:603765
1509	T	C P9	dbSNP:1065745
complement 1857	T	C P10	dbSNP:2301367
3565	G	C, A P11, P12	dbSNP:1065746
3594	T	G P13	dbSNP:1143646
3665	G	C P14	dbSNP:1065747
complement 4122	G	A P15	dbSNP:363099
complement 4985	G	A P16	dbSNP:363129
complement 5480	T	G P17	dbSNP:363125
6658	T	G P18	dbSNP:1143648
complement 6912	T	C P19	dbSNP:362336
complement 7753	G	A P20	dbSNP:3025816
complement 7849	G	C P21	dbSNP:3025814
complement 8478	T	C P22	dbSNP:2276881
8574	T	C P23	dbSNP:2229985
complement 9154	C	A P24	dbSNP:3025807
9498	T	C P25	dbSNP:2229987
complement 9699	G	A P26	dbSNP:362308
complement 9809	G	A P27	dbSNP:362307
complement 10064	T	C P28	dbSNP:362306
complement 10112	G	C P29	dbSNP:362268
complement 10124	G	C P30	dbSNP:362305
complement 10236	T	G P31	dbSNP:362304
complement 10271	G	A P32	dbSNP:362303
complement 10879	G	A P33	dbSNP:1557210
complement 10883	G	A P34	dbSNP:362302
complement 10971	C	A P35	dbSNP:3025805
complement 11181	G	A P36	dbSNP:362267
complement 11400	C	A P37	dbSNP:362301
11756..11757	G	- P38	dbSNP:5855774
12658	G	A P39	dbSNP:2237008
complement 12911	T	C P40	dbSNP:362300
complement 13040	G	A P41	dbSNP:2530595
13482	G	A P42	dbSNP:1803770
13563	G	A P43	dbSNP:1803771

Preferred *htt* SNPs have an allelic frequency of at least 30% in a sample population of patients. In preferred embodiments, a targeted *htt* SNP exhibits a frequency of heterozygosity of at least 25% within a sample patient population (*e.g.*, at least 30, 40, 50, 60, 65, 70, 75, or 80% heterozygosity).

In particularly preferred embodiment, the SNP allele is present at genomic site RS363125. In another particularly preferred embodiment, the SNP allele is present at genomic site RS362331. In another embodiment, the SNP allele is present at position 171, *e.g.*, an A171C polymorphism, in the huntingtin gene according to the sequence numbering in GenBank Accession No. NM_002111 (August 8, 2005).

II. anti-htt RNA Silencing Agents

The present invention features anti-huntingtin RNA silencing agents (*e.g.*, siRNA and shRNAs), methods of making said RNA silencing agents, and methods (*e.g.*, research and/or therapeutic methods) for using said improved RNA silencing agents (or portions thereof) for RNA silencing of mutant huntingtin protein. The RNA silencing agents comprise an antisense strand (or portions thereof), wherein the antisense strand has sufficient complementarity to a heterozygous single nucleotide polymorphism to mediate an RNA-mediated silencing mechanism (*e.g.* RNAi).

a) Design of anti-htt siRNA Molecules

An siRNA molecule of the invention is a duplex consisting of a sense strand and complementary antisense strand, the antisense strand having sufficient complementarity to a *htt* mRNA to mediate RNAi. Preferably, the siRNA molecule has a length from about 10-50 or more nucleotides, *i.e.*, each strand comprises 10-50 nucleotides (or nucleotide analogs). More preferably, the siRNA molecule has a length from about 16 - 30, *e.g.*, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, or 30 nucleotides in each strand, wherein one of the strands is sufficiently complementary to a target region. Preferably, the strands are aligned such that there are at least 1, 2, or 3 bases at the end of the strands which do not align (*i.e.*, for which no complementary bases occur in the opposing strand) such that an overhang of 1, 2 or 3 residues occurs at one or both ends of the duplex when strands are annealed. Preferably, the siRNA molecule has a length from about 10-50 or more nucleotides, *i.e.*, each strand comprises 10-50 nucleotides (or nucleotide analogs). More preferably, the siRNA molecule has a length from about 16 -

30, *e.g.*, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25, 26, 27, 28, 29, or 30 nucleotides in each strand, wherein one of the strands is substantially complementary to a target region *e.g.*, a gain-of-function gene target region, and the other strand is identical or substantially identical to the first strand.

Generally, siRNAs can be designed by using any method known in the art, for instance, by using the following protocol:

1. The siRNA should be specific for a heterozygous single-nucleotide polymorphism (SNP) found in a mutant huntingtin (htt) allele, but not a wild-type huntingtin allele. The first strand should be complementary to this sequence, and the other strand is substantially complementary to the first strand. In one embodiment, the SNP is outside the expanded CAG repeat of the mutant huntingtin (htt) allele. In another embodiment, the SNP is outside a coding region of the target gene. Exemplary polymorphisms are selected from the 5' untranslated region (5'-UTR) of a target gene. Cleavage of mRNA at these sites should eliminate translation of corresponding mutant protein. Polymorphisms from other regions of the mutant gene are also suitable for targeting. A sense strand is designed based on the sequence of the selected portion. Further, siRNAs with lower G/C content (35-55%) may be more active than those with G/C content higher than 55%. Thus in one embodiment, the invention includes nucleic acid molecules having 35-55% G/C content.

2. The sense strand of the siRNA is designed based on the sequence of the selected target site. Preferably the sense strand includes about 19 to 25 nucleotides, *e.g.*, 19, 20, 21, 22, 23, 24 or 25 nucleotides. More preferably, the sense strand includes 21, 22 or 23 nucleotides. The skilled artisan will appreciate, however, that siRNAs having a length of less than 19 nucleotides or greater than 25 nucleotides can also function to mediate RNAi. Accordingly, siRNAs of such length are also within the scope of the instant invention provided that they retain the ability to mediate RNAi. Longer RNA silencing agents have been demonstrated to elicit an interferon or PKR response in certain mammalian cells which may be undesirable. Preferably the RNA silencing agents of the invention do not elicit a PKR response (*i.e.*, are of a sufficiently short length). However, longer RNA silencing agents may be useful, for example, in cell

types incapable of generating a PRK response or in situations where the PKR response has been downregulated or dampened by alternative means.

The siRNA molecules of the invention have sufficient complementarity with the target site such that the siRNA can mediate RNAi. In general, siRNA containing nucleotide sequences sufficiently identical to a portion of the target gene to effect RISC-mediated cleavage of the target gene are preferred. Accordingly, in a preferred embodiment, the sense strand of the siRNA is designed have to have a sequence sufficiently identical to a portion of the target. For example, the sense strand may have 100% identity to the target site. However, 100% identity is not required. Greater than 80% identity, *e.g.*, 80%, 81%, 82%, 83%, 84%, 85%, 86%, 87%, 88%, 89%, 90%, 91%, 92%, 93%, 94%, 95%, 96%, 97%, 98%, 99% or even 100% identity, between the sense strand and the target RNA sequence is preferred. The invention has the advantage of being able to tolerate certain sequence variations to enhance efficiency and specificity of RNAi. In one embodiment, the sense strand has 4, 3, 2, 1, or 0 mismatched nucleotide(s) with a target region, such as a target region that differs by at least one base pair between the wild type and mutant allele, *e.g.*, a target region comprising the gain-of-function mutation, and the other strand is identical or substantially identical to the first strand. Moreover, siRNA sequences with small insertions or deletions of 1 or 2 nucleotides may also be effective for mediating RNAi. Alternatively, siRNA sequences with nucleotide analog substitutions or insertions can be effective for inhibition.

Sequence identity may determined by sequence comparison and alignment algorithms known in the art. To determine the percent identity of two nucleic acid sequences (or of two amino acid sequences), the sequences are aligned for optimal comparison purposes (*e.g.*, gaps can be introduced in the first sequence or second sequence for optimal alignment). The nucleotides (or amino acid residues) at corresponding nucleotide (or amino acid) positions are then compared. When a position in the first sequence is occupied by the same residue as the corresponding position in the second sequence, then the molecules are identical at that position. The percent identity between the two sequences is a function of the number of identical positions shared by the sequences (*i.e.*, % homology = # of identical positions/total # of positions x 100), optionally penalizing the score for the number of gaps introduced and/or length of gaps introduced.

The comparison of sequences and determination of percent identity between two sequences can be accomplished using a mathematical algorithm. In one embodiment, the alignment generated over a certain portion of the sequence aligned having sufficient identity but not over portions having low degree of identity (*i.e.*, a local alignment). A preferred, non-limiting example of a local alignment algorithm utilized for the comparison of sequences is the algorithm of Karlin and Altschul (1990) *Proc. Natl. Acad. Sci. USA* 87:2264-68, modified as in Karlin and Altschul (1993) *Proc. Natl. Acad. Sci. USA* 90:5873-77. Such an algorithm is incorporated into the BLAST programs (version 2.0) of Altschul, *et al.* (1990) *J. Mol. Biol.* 215:403-10.

In another embodiment, the alignment is optimized by introducing appropriate gaps and percent identity is determined over the length of the aligned sequences (*i.e.*, a gapped alignment). To obtain gapped alignments for comparison purposes, Gapped BLAST can be utilized as described in Altschul *et al.*, (1997) *Nucleic Acids Res.* 25(17):3389-3402. In another embodiment, the alignment is optimized by introducing appropriate gaps and percent identity is determined over the entire length of the sequences aligned (*i.e.*, a global alignment). A preferred, non-limiting example of a mathematical algorithm utilized for the global comparison of sequences is the algorithm of Myers and Miller, *CABIOS* (1989). Such an algorithm is incorporated into the ALIGN program (version 2.0) which is part of the GCG sequence alignment software package. When utilizing the ALIGN program for comparing amino acid sequences, a PAM120 weight residue table, a gap length penalty of 12, and a gap penalty of 4 can be used.

3. siRNAs are designed such that perfect complementarity exists between the siRNA and the target mRNA (*e.g.*, the mutant mRNA) at the polymorphism (*e.g.*, the point mutation), there thus being a mismatch if the siRNA is compared (*e.g.*, aligned) to the reference sequence (*e.g.*, wild type allele or mRNA sequence). The sense strand sequence may be designed such that the polymorphism is essentially in the middle of the strand. For example, if a 21-nucleotide siRNA is chosen, the polymorphism is at, for example, nucleotide 6, 7, 8, 9, 10, 11, 12, 13, 14, 15 or 16 (*i.e.*, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15 or 16 nucleotides from the 5' end of the sense strand. For a 22-nucleotide siRNA, the polymorphism is at, for example, nucleotide 7, 8, 9, 10, 11, 12, 13, 14, 15 or 16. For a 23-nucleotide siRNA, the polymorphism is at, for

example, 7, 8, 9, 10, 11, 12, 13, 14, 15 or 16. For a 24-nucleotide siRNA, the polymorphism is at, for example, 9, 10, 11, 12, 13, 14 or 16. For a 25-nucleotide siRNA, the polymorphism is at, for example, 9, 10, 11, 12, 13, 14, 15, 16 or 17.

In one preferred embodiment, the sense strand of the siRNA is identical to the polymorphism at a nucleotide position that is 10 nucleotides from the 5' end of the sense strand (*i.e.*, position P10).

In another preferred embodiment, the sense strand of the siRNA is identical to the polymorphism at a nucleotide position that is 16 nucleotides from the 5' end of the sense strand (*i.e.*, position P16).

4. siRNAs with single nucleotide specificity are preferably designed such that base pairing at the single nucleotide in the corresponding reference (*e.g.*, wild type) sequence is disfavored. For example, designing the siRNA such that purine:purine pairing exists between the siRNA and the wild type mRNA at the single nucleotide enhances single nucleotide specificity. The purine:purine pairing is selected, for example, from the group G:G, A:G, G:A and A:A pairing. Moreover, purine pyrimidine pairing between the siRNA and the mutant mRNA at the single nucleotide enhances single nucleotide specificity. The purine:pyrimidine pairing is selected, for example, from the group G:C, C:G, A:U, U:A, C:A, A:C, U:A and A:U pairing.

5. The antisense or guide strand of the siRNA is routinely the same length as the sense strand and includes complementary nucleotides. In one embodiment, the guide and sense strands are fully complementary, *i.e.*, the strands are blunt-ended when aligned or annealed. In another embodiment, the strands of the siRNA can be paired in such a way as to have a 3' overhang of 1 to 4, *e.g.*, 2, nucleotides. Overhangs can comprise (or consist of) nucleotides corresponding to the target gene sequence (or complement thereof). Alternatively, overhangs can comprise (or consist of) deoxyribonucleotides, for example dTs, or nucleotide analogs, or other suitable non-nucleotide material. Thus in another embodiment, the nucleic acid molecules may have a 3' overhang of 2 nucleotides, such as TT. The overhanging nucleotides may be either RNA or DNA. As noted above, it is desirable to choose a target region wherein the mutant:wild type mismatch is a purine:purine mismatch.

6. Using any method known in the art, compare the potential targets to the appropriate genome database (human, mouse, rat, etc.) and eliminate from consideration any target sequences with significant homology to other coding sequences. One such method for such sequence homology searches is known as BLAST, which is available at National Center for Biotechnology Information website.

7. Select one or more sequences that meet your criteria for evaluation.

Further general information about the design and use of siRNA may be found in "The siRNA User Guide," available at The Max-Planck-Institut für Biophysikalische Chemie website.

Alternatively, the siRNA may be defined functionally as a nucleotide sequence (or oligonucleotide sequence) that is capable of hybridizing with the target sequence (e.g., 400 mM NaCl, 40 mM PIPES pH 6.4, 1 mM EDTA, 50°C or 70°C hybridization for 12-16 hours; followed by washing). Additional preferred hybridization conditions include hybridization at 70°C in 1xSSC or 50°C in 1xSSC, 50% formamide followed by washing at 70°C in 0.3xSSC or hybridization at 70°C in 4xSSC or 50°C in 4xSSC, 50% formamide followed by washing at 67°C in 1xSSC. The hybridization temperature for hybrids anticipated to be less than 50 base pairs in length should be 5-10°C less than the melting temperature (T_m) of the hybrid, where T_m is determined according to the following equations. For hybrids less than 18 base pairs in length, $T_m(^{\circ}\text{C}) = 2(\# \text{ of A} + \text{T bases}) + 4(\# \text{ of G} + \text{C bases})$. For hybrids between 18 and 49 base pairs in length, $T_m(^{\circ}\text{C}) = 81.5 + 16.6(\log_{10}[\text{Na}^+]) + 0.41(\% \text{G} + \text{C}) - (600/\text{N})$, where N is the number of bases in the hybrid, and $[\text{Na}^+]$ is the concentration of sodium ions in the hybridization buffer ($[\text{Na}^+]$ for 1xSSC = 0.165 M). Additional examples of stringency conditions for polynucleotide hybridization are provided in Sambrook, J., E.F. Fritsch, and T. Maniatis, 1989, *Molecular Cloning: A Laboratory Manual*, Cold Spring Harbor Laboratory Press, Cold Spring Harbor, NY, chapters 9 and 11, and *Current Protocols in Molecular Biology*, 1995, F.M. Ausubel et al., eds., John Wiley & Sons, Inc., sections 2.10 and 6.3-6.4, incorporated herein by reference.

Negative control siRNAs should have the same nucleotide composition as the selected siRNA, but without significant sequence complementarity to the appropriate genome. Such negative controls may be designed by randomly scrambling the nucleotide sequence of the selected siRNA; a homology search can be performed to

ensure that the negative control lacks homology to any other gene in the appropriate genome. In addition, negative control siRNAs can be designed by introducing one or more base mismatches into the sequence.

8. To validate the effectiveness by which siRNAs destroy mutant mRNAs (e.g., mutant huntingtin mRNA), the siRNA may be incubated with mutant cDNA (e.g., mutant huntingtin cDNA) in a *Drosophila*-based *in vitro* mRNA expression system. Radiolabeled with ^{32}P , newly synthesized mutant mRNAs (e.g., mutant huntingtin mRNA) are detected autoradiographically on an agarose gel. The presence of cleaved mutant mRNA indicates mRNA nuclease activity. Suitable controls include omission of siRNA and use of wild-type huntingtin cDNA. Alternatively, control siRNAs are selected having the same nucleotide composition as the selected siRNA, but without significant sequence complementarity to the appropriate target gene. Such negative controls can be designed by randomly scrambling the nucleotide sequence of the selected siRNA; a homology search can be performed to ensure that the negative control lacks homology to any other gene in the appropriate genome. In addition, negative control siRNAs can be designed by introducing one or more base mismatches into the sequence.

ii. anti-htt siRNAs

Anti-*htt* siRNAs to target any of the single nucleotide polymorphisms described *supra*. Said siRNAs comprise an antisense strand which is fully complementary with the single nucleotide polymorphism. In certain embodiments, the RNA silencing agent is a siRNA.

In certain preferred embodiments, the siRNA comprises (i) a sense strand comprising the sequence set forth as SEQ ID NO: 3; and (ii) an antisense strand comprising the sequence set forth as SEQ ID NO: 4. In another embodiment, the siRNA comprises (i) a sense strand comprising the sequence set forth as SEQ ID NO: 7; and (ii) an antisense strand comprising the sequence set forth as SEQ ID NO: 8. In another embodiment, the siRNA comprises (i) a sense strand comprising the sequence set forth as SEQ ID NO: 9; and (ii) an antisense strand comprising the sequence set forth as SEQ ID NO: 10. In another embodiment, the siRNA comprises (i) a sense strand comprising the sequence set forth as SEQ ID NO: 13; and (ii) an antisense strand comprising the

sequence set forth as SEQ ID NO: 14. In another embodiment, the siRNA comprises (i) a sense strand comprising the sequence set forth as SEQ ID NO: 15; and (ii) an antisense strand comprising the sequence set forth as SEQ ID NO: 16. In another embodiment, the siRNA comprises (i) a sense strand comprising the sequence set forth as SEQ ID NO: 18; and (ii) an antisense strand comprising the sequence set forth as SEQ ID NO: 19.

To validate the effectiveness by which siRNAs destroy mutant mRNAs (e.g., mutant huntingtin mRNA), the siRNA is incubated with mutant cDNA (e.g., mutant huntingtin cDNA) in a *Drosophila*-based *in vitro* mRNA expression system. Radiolabeled with ^{32}P , newly synthesized mutant mRNAs (e.g., mutant huntingtin mRNA) are detected autoradiographically on an agarose gel. The presence of cleaved mutant mRNA indicates mRNA nuclease activity. Suitable controls include omission of siRNA and use of wild-type huntingtin cDNA. Alternatively, control siRNAs are selected having the same nucleotide composition as the selected siRNA, but without significant sequence complementarity to the appropriate target gene. Such negative controls can be designed by randomly scrambling the nucleotide sequence of the selected siRNA; a homology search can be performed to ensure that the negative control lacks homology to any other gene in the appropriate genome. In addition, negative control siRNAs can be designed by introducing one or more base mismatches into the sequence.

Sites of siRNA-mRNA complementation are selected which result in optimal mRNA specificity and maximal mRNA cleavage.

While the instant invention primarily features targeting polymorphic regions in the target mutant gene (e.g., in mutant htt) distinct from the expanded CAG region mutation, the skilled artisan will appreciate that targeting the mutant region may have applicability as a therapeutic strategy in certain situations. Targeting the mutant region can be accomplished using siRNA that complements CAG in series. The siRNA^{CAG} would bind to mRNAs with CAG complementation, but might be expected to have greater opportunity to bind to an extended CAG series. Multiple siRNA^{CAG} would bind to the mutant huntingtin mRNA (as opposed to fewer for the wild type huntingtin mRNA); thus, the mutant huntingtin mRNA is more likely to be cleaved. Successful mRNA inactivation using this approach would also eliminate normal or wild-type

huntingtin mRNA. Also inactivated, at least to some extent, could be other normal genes (approximately 70) which also have CAG repeats, where their mRNAs could interact with the siRNA. This approach would thus rely on an attrition strategy -- more of the mutant huntingtin mRNA would be destroyed than wild type huntingtin mRNA or the other approximately 69 mRNAs that code for polyglutamines.

b) siRNA-like molecules

siRNA-like molecules of the invention have a sequence (i.e., have a strand having a sequence) that is "sufficiently complementary" to a heterozygous SNP of a htt mRNA to direct gene silencing either by RNAi or translational repression. siRNA-like molecules are designed in the same way as siRNA molecules, but the degree of sequence identity between the sense strand and target RNA approximates that observed between an miRNA and its target. In general, as the degree of sequence identity between a miRNA sequence and the corresponding target gene sequence is decreased, the tendency to mediate post-transcriptional gene silencing by translational repression rather than RNAi is increased. Therefore, in an alternative embodiment, where post-transcriptional gene silencing by translational repression of the target gene is desired, the miRNA sequence has partial complementarity with the target gene sequence. In certain embodiments, the miRNA sequence has partial complementarity with one or more short sequences (complementarity sites) dispersed within the target mRNA (e.g. within the 3'-UTR of the target mRNA) (Hutvagner and Zamore, Science, 2002; Zeng et al., Mol. Cell, 2002; Zeng et al., RNA, 2003; Doench et al., Genes & Dev., 2003). Since the mechanism of translational repression is cooperative, multiple complementarity sites (e.g., 2, 3, 4, 5, or 6) may be targeted in certain embodiments.

The capacity of a siRNA-like duplex to mediate RNAi or translational repression may be predicted by the distribution of non-identical nucleotides between the target gene sequence and the nucleotide sequence of the silencing agent at the site of complementarity. In one embodiment, where where gene silencing by translational repression is desired, at least one non-identical nucleotide is present in the central portion of the complementarity site so that duplex formed by the miRNA guide strand and the target mRNA contains a central "bulge" (Doench JG et al., Genes & Dev., 2003). In another embodiment 2, 3, 4, 5, or 6 contiguous or non-contiguous non-

identical nucleotides are introduced. The non-identical nucleotide may be selected such that it forms a wobble base pair (e.g., G:U) or a mismatched base pair (G:A, C:A, C:U, G:G, A:A, C:C, U:U). In a further preferred embodiment, the "bulge" is centered at nucleotide positions 12 and 13 from the 5' end of the miRNA molecule.

c) Short hairpin RNA (shRNA) molecules

In certain featured embodiments, the instant invention provides shRNAs capable of mediating RNA silencing of a heterozygous htt SNP with enhanced selectivity. In contrast to siRNAs, shRNAs mimic the natural precursors of micro RNAs (miRNAs) and enter at the top of the gene silencing pathway. For this reason, shRNAs are believed to mediate gene silencing more efficiently by being fed through the entire natural gene silencing pathway.

miRNAs are noncoding RNAs of approximately 22 nucleotides which can regulate gene expression at the post transcriptional or translational level during plant and animal development. One common feature of miRNAs is that they are all excised from an approximately 70 nucleotide precursor RNA stem-loop termed pre-miRNA, probably by Dicer, an RNase III-type enzyme, or a homolog thereof. Naturally-occurring miRNA precursors (pre-miRNA) have a single strand that forms a duplex stem including two portions that are generally complementary, and a loop, that connects the two portions of the stem. In typical pre-miRNAs, the stem includes one or more bulges, e.g., extra nucleotides that create a single nucleotide "loop" in one portion of the stem, and/or one or more unpaired nucleotides that create a gap in the hybridization of the two portions of the stem to each other. Short hairpin RNAs, or engineered RNA precursors, of the invention are artificial constructs based on these naturally occurring pre-miRNAs, but which are engineered to deliver desired RNA silencing agents (e.g., siRNAs of the invention). By substituting the stem sequences of the pre-miRNA with sequence complementary to the target mRNA, a shRNA is formed. The shRNA is processed by the entire gene silencing pathway of the cell, thereby efficiently mediating RNAi.

The requisite elements of a shRNA molecule include a first portion and a second portion, having sufficient complementarity to anneal or hybridize to form a duplex or double-stranded stem portion. The two portions need not be fully or perfectly

complementary. The first and second "stem" portions are connected by a portion having a sequence that has insufficient sequence complementarity to anneal or hybridize to other portions of the shRNA. This latter portion is referred to as a "loop" portion in the shRNA molecule. The shRNA molecules are processed to generate siRNAs. shRNAs can also include one or more bulges, *i.e.*, extra nucleotides that create a small nucleotide "loop" in a portion of the stem, for example a one-, two- or three-nucleotide loop. The stem portions can be the same length, or one portion can include an overhang of, for example, 1-5 nucleotides. The overhanging nucleotides can include, for example, uracils (Us), *e.g.*, all Us. Such Us are notably encoded by thymidines (Ts) in the shRNA-encoding DNA which signal the termination of transcription.

In shRNAs, or engineered precursor RNAs, of the instant invention, one portion of the duplex stem is a nucleic acid sequence that is complementary (or anti-sense) to the heterozygous SNP. Preferably, one strand of the stem portion of the shRNA is sufficiently complementary (*e.g.*, antisense) to a target RNA (*e.g.*, mRNA) sequence to mediate degradation or cleavage of said target RNA *via* RNA interference (RNAi). Thus, engineered RNA precursors include a duplex stem with two portions and a loop connecting the two stem portions. The antisense portion can be on the 5' or 3' end of the stem. The stem portions of a shRNA are preferably about 15 to about 50 nucleotides in length. Preferably the two stem portions are about 18 or 19 to about 21, 22, 23, 24, 25, 30, 35, 37, 38, 39, or 40 or more nucleotides in length. In preferred embodiments, the length of the stem portions should be 21 nucleotides or greater. When used in mammalian cells, the length of the stem portions should be less than about 30 nucleotides to avoid provoking non-specific responses like the interferon pathway. In non-mammalian cells, the stem can be longer than 30 nucleotides. In fact, the stem can include much larger sections complementary to the target mRNA (up to, and including the entire mRNA). In fact, a stem portion can include much larger sections complementary to the target mRNA (up to, and including the entire mRNA).

The two portions of the duplex stem must be sufficiently complementary to hybridize to form the duplex stem. Thus, the two portions can be, but need not be, fully or perfectly complementary. In addition, the two stem portions can be the same length, or one portion can include an overhang of 1, 2, 3, or 4 nucleotides. The overhanging nucleotides can include, for example, uracils (Us), *e.g.*, all Us. The loop in the shRNAs or engineered RNA precursors may differ from natural pre-miRNA sequences by

modifying the loop sequence to increase or decrease the number of paired nucleotides, or replacing all or part of the loop sequence with a tetraloop or other loop sequences. Thus, the loop in the shRNAs or engineered RNA precursors can be 2, 3, 4, 5, 6, 7, 8, 9, or more, *e.g.*, 15 or 20, or more nucleotides in length.

The loop in the shRNAs or engineered RNA precursors may differ from natural pre-miRNA sequences by modifying the loop sequence to increase or decrease the number of paired nucleotides, or replacing all or part of the loop sequence with a tetraloop or other loop sequences. Thus, the loop portion in the shRNA can be about 2 to about 20 nucleotides in length, *i.e.*, about 2, 3, 4, 5, 6, 7, 8, 9, or more, *e.g.*, 15 or 20, or more nucleotides in length. A preferred loop consists of or comprises a "tetraloop" sequences. Exemplary tetraloop sequences include, but are not limited to, the sequences GNRA, where N is any nucleotide and R is a purine nucleotide, GGGG, and UUUU.

In certain embodiments, shRNAs of the invention include the sequences of a desired siRNA molecule described *supra*. In other embodiments, the sequence of the antisense portion of a shRNA can be designed essentially as described above or generally by selecting an 18, 19, 20, 21 nucleotide, or longer, sequence from within the target RNA (*e.g.*, SOD1 or *hlt* mRNA), for example, from a region 100 to 200 or 300 nucleotides upstream or downstream of the start of translation. In general, the sequence can be selected from any portion of the target RNA (*e.g.*, mRNA) including the 5' UTR (untranslated region), coding sequence, or 3' UTR, provided said portion is distant from the site of the gain-of-function mutation. This sequence can optionally follow immediately after a region of the target gene containing two adjacent AA nucleotides. The last two nucleotides of the nucleotide sequence can be selected to be UU. This 21 or so nucleotide sequence is used to create one portion of a duplex stem in the shRNA. This sequence can replace a stem portion of a wild-type pre-miRNA sequence, *e.g.*, enzymatically, or is included in a complete sequence that is synthesized. For example, one can synthesize DNA oligonucleotides that encode the entire stem-loop engineered RNA precursor, or that encode just the portion to be inserted into the duplex stem of the precursor, and using restriction enzymes to build the engineered RNA precursor construct, *e.g.*, from a wild-type pre-miRNA.

Engineered RNA precursors include in the duplex stem the 21-22 or so nucleotide sequences of the siRNA or siRNA-like duplex desired to be produced *in vivo*. Thus, the stem portion of the engineered RNA precursor includes at least 18 or 19

nucleotide pairs corresponding to the sequence of an exonic portion of the gene whose expression is to be reduced or inhibited. The two 3' nucleotides flanking this region of the stem are chosen so as to maximize the production of the siRNA from the engineered RNA precursor and to maximize the efficacy of the resulting siRNA in targeting the corresponding mRNA for translational repression or destruction by RNAi *in vivo* and *in vitro*.

In certain embodiments, shRNAs of the invention include miRNA sequences, optionally end-modified miRNA sequences, to enhance entry into RISC. The miRNA sequence can be similar or identical to that of any naturally occurring miRNA (see *e.g.* The miRNA Registry; Griffiths-Jones S, *Nuc. Acids Res.*, 2004). Over one thousand natural miRNAs have been identified to date and together they are thought to comprise ~1% of all predicted genes in the genome. Many natural miRNAs are clustered together in the introns of pre-mRNAs and can be identified *in silico* using homology-based searches (Pasquinelli et al., 2000; Lagos-Quintana et al., 2001; Lau et al., 2001; Lee and Ambros, 2001) or computer algorithms (*e.g.* MiRScan, MiRSeeker) that predict the capability of a candidate miRNA gene to form the stem loop structure of a pri-mRNA (Grad et al., *Mol. Cell.*, 2003; Lim et al., *Genes Dev.*, 2003; Lim et al., *Science*, 2003; Lai EC et al., *Genome Bio.*, 2003). An online registry provides a searchable database of all published miRNA sequences (The miRNA Registry at the Sanger Institute website; Griffiths-Jones S, *Nuc. Acids Res.*, 2004). Exemplary, natural miRNAs include lin-4, let-7, miR-10, mirR-15, miR-16, miR-168, miR-175, miR-196 and their homologs, as well as other natural miRNAs from humans and certain model organisms including *Drosophila melanogaster*, *Caenorhabditis elegans*, zebrafish, *Arabidopsis thaliana*, mouse, and rat as described in International PCT Publication No. WO 03/029459.

Naturally-occurring miRNAs are expressed by endogenous genes *in vivo* and are processed from a hairpin or stem-loop precursor (pre-miRNA or pri-miRNAs) by Dicer or other RNases (Lagos-Quintana et al., *Science*, 2001; Lau et al., *Science*, 2001; Lee and Ambros, *Science*, 2001; Lagos-Quintana et al., *Curr. Biol.*, 2002; Mourelatos et al., *Genes Dev.*, 2002; Reinhart et al., *Science*, 2002; Ambros et al., *Curr. Biol.*, 2003; Brennecke et al., 2003; Lagos-Quintana et al., *RNA*, 2003; Lim et al., *Genes Dev.*, 2003; Lim et al., *Science*, 2003). miRNAs can exist transiently *in vivo* as a double-stranded duplex but only one strand is taken up by the RISC complex to direct gene silencing.

Certain miRNAs, *e.g.* plant miRNAs, have perfect or near-perfect complementarity to their target mRNAs and, hence, direct cleavage of the target mRNAs. Other miRNAs have less than perfect complementarity to their target mRNAs and, hence, direct translational repression of the target mRNAs. The degree of complementarity between an miRNA and its target mRNA is believed to determine its mechanism of action. For example, perfect or near-perfect complementarity between a miRNA and its target mRNA is predictive of a cleavage mechanism (Yekta *et al.*, *Science*, 2004), whereas less than perfect complementarity is predictive of a translational repression mechanism. In particular embodiments, the miRNA sequence is that of a naturally-occurring miRNA sequence, the aberrant expression or activity of which is correlated with a miRNA disorder.

d) Dual Functional Oligonucleotide Tethers

In other embodiments, the RNA silencing agents of the present invention include dual functional oligonucleotide tethers useful for the intercellular recruitment of a miRNA. Animal cells express a range of miRNAs, noncoding RNAs of approximately 22 nucleotides which can regulate gene expression at the post transcriptional or translational level. By binding a miRNA bound to RISC and recruiting it to a target mRNA, a dual functional oligonucleotide tether can repress the expression of genes involved *e.g.*, in the arteriosclerotic process. The use of oligonucleotide tethers offer several advantages over existing techniques to repress the expression of a particular gene. First, the methods described herein allow an endogenous molecule (often present in abundance), an miRNA, to mediate RNA silencing; accordingly the methods described herein obviate the need to introduce foreign molecules (*e.g.*, siRNAs) to mediate RNA silencing. Second, the RNA-silencing agents and, in particular, the linking moiety (*e.g.*, oligonucleotides such as the 2'-*O*-methyl oligonucleotide), can be made stable and resistant to nuclease activity. As a result, the tethers of the present invention can be designed for direct delivery, obviating the need for indirect delivery (*e.g.* viral) of a precursor molecule or plasmid designed to make the desired agent within the cell. Third, tethers and their respective moieties, can be designed to conform to specific mRNA sites and specific miRNAs. The designs can be cell and gene product specific. Fourth, the methods disclosed herein leave the mRNA intact, allowing one

skilled in the art to block protein synthesis in short pulses using the cell's own machinery. As a result, these methods of RNA silencing are highly regulatable.

The dual functional oligonucleotide tethers ("tethers") of the invention are designed such that they recruit miRNAs (*e.g.*, endogenous cellular miRNAs) to a target mRNA so as to induce the modulation of a gene of interest. In preferred embodiments, the tethers have the formula T-L- μ , wherein T is an mRNA targeting moiety, L is a linking moiety, and μ is an miRNA recruiting moiety. Any one or more moiety may be double stranded. Preferably, however, each moiety is single stranded.

Moieties within the tethers can be arranged or linked (in the 5' to 3' direction) as depicted in the formula T-L- μ (*i.e.*, the 3' end of the targeting moiety linked to the 5' end of the linking moiety and the 3' end of the linking moiety linked to the 5' end of the miRNA recruiting moiety). Alternatively, the moieties can be arranged or linked in the tether as follows: μ -T-L (*i.e.*, the 3' end of the miRNA recruiting moiety linked to the 5' end of the linking moiety and the 3' end of the linking moiety linked to the 5' end of the targeting moiety).

The mRNA targeting moiety, as described above, is capable of capturing a specific target mRNA. According to the invention, expression of the target mRNA is undesirable, and, thus, translational repression of the mRNA is desired. The mRNA targeting moiety should be of sufficient size to effectively bind the target mRNA. The length of the targeting moiety will vary greatly depending, in part, on the length of the target mRNA and the degree of complementarity between the target mRNA and the targeting moiety. In various embodiments, the targeting moiety is less than about 200, 100, 50, 30, 25, 20, 19, 18, 17, 16, 15, 14, 13, 12, 11, 10, 9, 8, 7, 6, or 5 nucleotides in length. In a particular embodiment, the targeting moiety is about 15 to about 25 nucleotides in length.

The miRNA recruiting moiety, as described above, is capable of associating with a miRNA. According to the invention, the miRNA may be any miRNA capable of repressing the target mRNA. Mammals are reported to have over 250 endogenous miRNAs (Lagos-Quintana *et al.* (2002) *Current Biol.* 12:735-739; Lagos-Quintana *et al.* (2001) *Science* 294:858-862; and Lim *et al.* (2003) *Science* 299:1540). In various embodiments, the miRNA may be any art-recognized miRNA. Table 3 lists some of the known human miRNAs

The linking moiety is any agent capable of linking the targeting moieties such that the activity of the targeting moieties is maintained. Linking moieties are preferably oligonucleotide moieties comprising a sufficient number of nucleotides such that the targeting agents can sufficiently interact with their respective targets. Linking moieties have little or no sequence homology with cellular mRNA or miRNA sequences. Exemplary linking moieties include one or more 2'-O- methylnucleotides, *e.g.*, 2'-O-methyladenosine, 2'-O-methylthymidine, 2'-O-methylguanosine or 2'-O-methyluridine.

III. Modified anti-*htr* RNA Silencing Agents

In certain aspects of the invention, an RNA silencing agent (or any portion thereof) of the invention as described *supra* may be modified such that the activity of the agent is further improved. For example, the RNA silencing agents described in Section II *supra* may be modified with any of the modifications described *infra*. The modifications can, in part, serve to further enhance target discrimination, to enhance stability of the agent (*e.g.*, to prevent degradation), to promote cellular uptake, to enhance the target efficiency, to improve efficacy in binding (*e.g.*, to the targets), to improve patient tolerance to the agent, and/or to reduce toxicity.

1) *Modifications to Enhance Target Discrimination*

In certain embodiments, the RNA silencing agents of the invention may be substituted with a destabilizing nucleotide to enhance single nucleotide target discrimination (see US Application No. 11/698,689, filed January 25, 2007 and US Provisional Application No. 60/762,225 filed January 25, 2006, both of which are incorporated herein by reference). Such a modification may be sufficient to abolish the specificity of the RNA silencing agent for a non-target mRNA (*e.g.* wild-type mRNA), without appreciably affecting the specificity of the RNA silencing agent for a target mRNA (*e.g.* gain-of-function mutant mRNA).

In preferred embodiments, the RNA silencing agents of the invention are modified by the introduction of at least one universal nucleotide in the antisense strand thereof. Universal nucleotides comprise base portions that are capable of base pairing indiscriminately with any of the four conventional nucleotide bases (*e.g.* A,G,C,U). A universal nucleotide is preferred because it has relatively minor effect on the stability of the RNA duplex or the duplex formed by the guide strand of the RNA silencing agent

and the target mRNA. Exemplary universal nucleotide include those having an inosine base portion or an inosine analog base portion selected from the group consisting of deoxyinosine (e.g. 2'-deoxyinosine), 7-deaza-2'-deoxyinosine, 2'-aza-2'-deoxyinosine, PNA-inosine, morpholino-inosine, LNA-inosine, phosphoramidate-inosine, 2'-O-methoxyethyl-inosine, and 2'-OMe-inosine. In particularly preferred embodiments, the universal nucleotide is an inosine residue or a naturally occurring analog thereof.

The RNA silencing agents of the invention are preferably modified by the introduction of at least one destabilizing nucleotide within 5 nucleotides from a specificity-determining nucleotide (*ie.* the nucleotide which recognizes the disease-related polymorphism). For example, the destabilizing nucleotide may be introduced at a position that is within 5, 4, 3, 2, or 1 nucleotide(s) from a specificity-determining nucleotide. In exemplary embodiments, the destabilizing nucleotide is introduced at a position which is 3 nucleotides from the specificity-determining nucleotide (*ie.* such that there are 2 stabilizing nucleotides between the destabilizing nucleotide and the specificity-determining nucleotide). In RNA silencing agents having two strands or strand portions (e.g. siRNAs and shRNAs), the destabilizing nucleotide may be introduced in the strand or strand portion that does not contain the specificity-determining nucleotide. In preferred embodiments, the destabilizing nucleotide is introduced in the same strand or strand portion that contains the specificity-determining nucleotide.

2) Modifications to Enhance Efficacy and Specificity

In certain embodiments, the RNA silencing agents of the invention may be altered to facilitate enhanced efficacy and specificity in mediating RNAi according to asymmetry design rules (see International Publication No. WO 2005/001045, US Publication No. 2005-0181382 A1). Such alterations facilitate entry of the antisense strand of the siRNA (e.g., a siRNA designed using the methods of the invention or an siRNA produced from a shRNA) into RISC in favor of the sense strand, such that the antisense strand preferentially guides cleavage or translational repression of a target mRNA, and thus increasing or improving the efficiency of target cleavage and silencing. Preferably the asymmetry of an RNA silencing agent is enhanced by lessening the base pair strength between the antisense strand 5' end (AS 5') and the sense strand 3' end (S 3') of the RNA silencing agent relative to the bond strength or base pair strength

between the antisense strand 3' end (AS 3') and the sense strand 5' end (S '5) of said RNA silencing agent.

In one embodiment, the asymmetry of an RNA silencing agent of the invention may be enhanced such that there are fewer G:C base pairs between the 5' end of the first or antisense strand and the 3' end of the sense strand portion than between the 3' end of the first or antisense strand and the 5' end of the sense strand portion. In another embodiment, the asymmetry of an RNA silencing agent of the invention may be enhanced such that there is at least one mismatched base pair between the 5' end of the first or antisense strand and the 3' end of the sense strand portion. Preferably, the mismatched base pair is selected from the group consisting of G:A, C:A, C:U, G:G, A:A, C:C and U:U. In another embodiment, the asymmetry of an RNA silencing agent of the invention may be enhanced such that there is at least one wobble base pair, e.g., G:U, between the 5' end of the first or antisense strand and the 3' end of the sense strand portion. In another embodiment, the asymmetry of an RNA silencing agent of the invention may be enhanced such that there is at least one base pair comprising a rare nucleotide, e.g., inosine (I). Preferably, the base pair is selected from the group consisting of an I:A, I:U and I:C. In yet another embodiment, the the asymmetry of an RNA silencing agent of the invention may be enhanced such that there is at least one base pair comprising a modified nucleotide. In preferred embodiments, the modified nucleotide is selected from the group consisting of 2-amino-G, 2-amino-A, 2,6-diamino-G, and 2,6-diamino-A.

3) *RNA silencing agents with Enhanced Stability*

The RNA silencing agents of the present invention can be modified to improve stability in serum or in growth medium for cell cultures. In order to enhance the stability, the 3'-residues may be stabilized against degradation, e.g., they may be selected such that they consist of purine nucleotides, particularly adenosine or guanosine nucleotides. Alternatively, substitution of pyrimidine nucleotides by modified analogues, e.g., substitution of uridine by 2'-deoxythymidine is tolerated and does not affect the efficiency of RNA interference.

In a preferred aspect, the invention features RNA silencing agents that include first and second strands wherein the second strand and/or first strand is modified by the substitution of internal nucleotides with modified nucleotides, such that *in vivo* stability

is enhanced as compared to a corresponding unmodified RNA silencing agent. As defined herein, an "internal" nucleotide is one occurring at any position other than the 5' end or 3' end of nucleic acid molecule, polynucleotide or oligonucleotide. An internal nucleotide can be within a single-stranded molecule or within a strand of a duplex or double-stranded molecule. In one embodiment, the sense strand and/or antisense strand is modified by the substitution of at least one internal nucleotide. In another embodiment, the sense strand and/or antisense strand is modified by the substitution of at least 2, 3, 4, 5, 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 21, 22, 23, 24, 25 or more internal nucleotides. In another embodiment, the sense strand and/or antisense strand is modified by the substitution of at least 5%, 10%, 15%, 20%, 25%, 30%, 35%, 40%, 45%, 50%, 55%, 60%, 65%, 70%, 75%, 80%, 85%, 90%, 95% or more of the internal nucleotides. In yet another embodiment, the sense strand and/or antisense strand is modified by the substitution of all of the internal nucleotides.

In a preferred embodiment of the present invention the RNA silencing agents may contain at least one modified nucleotide analogue. The nucleotide analogues may be located at positions where the target-specific silencing activity, *e.g.*, the RNAi mediating activity or translational repression activity is not substantially effected, *e.g.*, in a region at the 5'-end and/or the 3'-end of the siRNA molecule. Particularly, the ends may be stabilized by incorporating modified nucleotide analogues.

Preferred nucleotide analogues include sugar- and/or backbone-modified ribonucleotides (*i.e.*, include modifications to the phosphate-sugar backbone). For example, the phosphodiester linkages of natural RNA may be modified to include at least one of a nitrogen or sulfur heteroatom. In preferred backbone-modified ribonucleotides the phosphoester group connecting to adjacent ribonucleotides is replaced by a modified group, *e.g.*, of phosphothioate group. In preferred sugar-modified ribonucleotides, the 2' OH-group is replaced by a group selected from H, OR, R, halo, SH, SR, NH₂, NHR, NR₂ or ON, wherein R is C₁-C₆ alkyl, alkenyl or alkynyl and halo is F, Cl, Br or I.

In particular embodiments, the modifications are 2'-fluoro, 2'-amino and/or 2'-thio modifications. Particularly preferred modifications include 2'-fluoro-cytidine, 2'-fluoro-uridine, 2'-fluoro-adenosine, 2'-fluoro-guanosine, 2'-amino-cytidine, 2'-amino-uridine, 2'-amino-adenosine, 2'-amino-guanosine, 2,6-diaminopurine, 4-thio-uridine, and/or 5-amino-allyl-uridine. In a particular embodiment, the 2'-fluoro ribonucleotides

are every uridine and cytidine. Additional exemplary modifications include 5-bromo-uridine, 5-iodo-uridine, 5-methyl-cytidine, ribo-thymidine, 2-aminopurine, 2'-amino-butyl-uridine, 5-fluoro-cytidine, and 5-fluoro-uridine. 2'-deoxy-nucleotides and 2'-Ome nucleotides can also be used within modified RNA-silencing agents moieties of the instant invention. Additional modified residues include, deoxy-abasic, inosine, N3-methyl-uridine, N6, N6-dimethyl-adenosine, pseudouridine, purine ribonucleoside and ribavirin. In a particularly preferred embodiment, the 2' moiety is a methyl group such that the linking moiety is a 2'-O-methyl oligonucleotide.

In an exemplary embodiment, the RNA silencing agent of the invention comprises Locked Nucleic Acids (LNAs). LNAs comprise sugar-modified nucleotides that resist nuclease activities (are highly stable) and possess single nucleotide discrimination for mRNA (Elmen *et al.*, *Nucleic Acids Res.*, (2005), 33(1): 439-447; Braasch *et al.* (2003) *Biochemistry* 42:7967-7975, Petersen *et al.* (2003) *Trends Biotechnol* 21:74-81). These molecules have 2'-O,4'-C-ethylene-bridged nucleic acids, with possible modifications such as 2'-deoxy-2"-fluorouridine. Moreover, LNAs increase the specificity of oligonucleotides by constraining the sugar moiety into the 3'-endo conformation, thereby preorganizing the nucleotide for base pairing and increasing the melting temperature of the oligonucleotide by as much as 10°C per base.

In another exemplary embodiment, the RNA silencing agent of the invention comprises Peptide Nucleic Acids (PNAs). PNAs comprise modified nucleotides in which the sugar-phosphate portion of the nucleotide is replaced with a neutral 2-amino ethylglycine moiety capable of forming a polyamide backbone which is highly resistant to nuclease digestion and imparts improved binding specificity to the molecule (Nielsen, *et al.*, *Science*, (2001), 254: 1497-1500).

Also preferred are nucleobase-modified ribonucleotides, *i.e.*, ribonucleotides, containing at least one non-naturally occurring nucleobase instead of a naturally occurring nucleobase. Bases may be modified to block the activity of adenosine deaminase. Exemplary modified nucleobases include, but are not limited to, uridine and/or cytidine modified at the 5-position, *e.g.*, 5-(2-amino)propyl uridine, 5-bromo uridine; adenosine and/or guanosines modified at the 8 position, *e.g.*, 8-bromo guanosine; deaza nucleotides, *e.g.*, 7-deaza-adenosine; O- and N-alkylated nucleotides, *e.g.*, N6-methyl adenosine are suitable. It should be noted that the above modifications may be combined.

In other embodiments, cross-linking can be employed to alter the pharmacokinetics of the RNA silencing agent, for example, to increase half-life in the body. Thus, the invention includes RNA silencing agents having two complementary strands of nucleic acid, wherein the two strands are crosslinked. The invention also includes RNA silencing agents which are conjugated or unconjugated (*e.g.*, at its 3' terminus) to another moiety (*e.g.* a non-nucleic acid moiety such as a peptide), an organic compound (*e.g.*, a dye), or the like). Modifying siRNA derivatives in this way may improve cellular uptake or enhance cellular targeting activities of the resulting siRNA derivative as compared to the corresponding siRNA, are useful for tracing the siRNA derivative in the cell, or improve the stability of the siRNA derivative compared to the corresponding siRNA.

Other exemplary modifications include: (a) 2' modification, *e.g.*, provision of a 2' OMe moiety on a U in a sense or antisense strand, but especially on a sense strand, or provision of a 2' OMe moiety in a 3' overhang, *e.g.*, at the 3' terminus (3' terminus means at the 3' atom of the molecule or at the most 3' moiety, *e.g.*, the most 3' P or 2' position, as indicated by the context); (b) modification of the backbone, *e.g.*, with the replacement of an O with an S, in the phosphate backbone, *e.g.*, the provision of a phosphorothioate modification, on the U or the A or both, especially on an antisense strand; *e.g.*, with the replacement of a P with an S; (c) replacement of the U with a C5 amino linker; (d) replacement of an A with a G (sequence changes are preferred to be located on the sense strand and not the antisense strand); and (d) modification at the 2', 6', 7', or 8' position. Preferred embodiments are those in which one or more of these modifications are present on the sense but not the antisense strand, or embodiments where the antisense strand has fewer of such modifications. Yet other exemplary modifications include the use of a methylated P in a 3' overhang, *e.g.*, at the 3' terminus; combination of a 2' modification, *e.g.*, provision of a 2' O Me moiety and modification of the backbone, *e.g.*, with the replacement of a P with an S, *e.g.*, the provision of a phosphorothioate modification, or the use of a methylated P, in a 3' overhang, *e.g.*, at the 3' terminus; modification with a 3' alkyl; modification with an abasic pyrrolidine in a 3' overhang, *e.g.*, at the 3' terminus; modification with naproxen, ibuprofen, or other moieties which inhibit degradation at the 3' terminus.

4) *Modifications to Enhance Cellular Uptake*

In other embodiments, RNA silencing agents may be modified with chemical moieties, for example, to enhance cellular uptake by target cells (*e.g.*, neuronal cells). Thus, the invention includes RNA silencing agents which are conjugated or unconjugated (*e.g.*, at its 3' terminus) to another moiety (*e.g.* a non-nucleic acid moiety such as a peptide), an organic compound (*e.g.*, a dye), or the like. The conjugation can be accomplished by methods known in the art, *e.g.*, using the methods of Lambert *et al.*, *Drug Deliv. Rev.* 47(1), 99-112 (2001) (describes nucleic acids loaded to polyalkylcyanoacrylate (PACA) nanoparticles); Fattal *et al.*, *J. Control Release* 53(1-3):137-43 (1998) (describes nucleic acids bound to nanoparticles); Schwab *et al.*, *Ann. Oncol.* 5 Suppl. 4:55-8 (1994) (describes nucleic acids linked to intercalating agents, hydrophobic groups, polycations or PACA nanoparticles); and Godard *et al.*, *Eur. J. Biochem.* 232(2):404-10 (1995) (describes nucleic acids linked to nanoparticles).

In a particular embodiment, an RNA silencing agent of invention is conjugated to a lipophilic moiety. In one embodiment, the lipophilic moiety is a ligand that includes a cationic group. In another embodiment, the lipophilic moiety is attached to one or both strands of an siRNA. In a preferred embodiment, the lipophilic moiety is attached to one end of the sense strand of the siRNA. In another preferred embodiment, the lipophilic moiety is attached to the 3' end of the sense strand. In certain embodiments, the lipophilic moiety is selected from the group consisting of cholesterol, vitamin E, vitamin K, vitamin A, folic acid, or a cationic dye (*e.g.*, Cy3). In a preferred embodiment, the lipophilic moiety is a cholesterol. Other lipophilic moieties include cholic acid, adamantane acetic acid, 1-pyrene butyric acid, dihydrotestosterone, 1,3-Bis-O(hexadecyl)glycerol, geranyloxyhexyl group, hexadecylglycerol, borneol, menthol, 1,3-propanediol, heptadecyl group, palmitic acid, myristic acid, O3-(oleoyl)lithocholic acid, O3-(oleoyl)cholenic acid, dimethoxytrityl, or phenoxazine.

5) *Tethered Ligands*

Other entities can be tethered to an RNA silencing agent of the invention. For example, a ligand tethered to an RNA silencing agent to improve stability, hybridization thermodynamics with a target nucleic acid, targeting to a particular tissue or cell-type, or cell permeability, *e.g.*, by an endocytosis-dependent or -independent mechanism.

Ligands and associated modifications can also increase sequence specificity and consequently decrease off-site targeting. A tethered ligand can include one or more modified bases or sugars that can function as intercalators. These are preferably located in an internal region, such as in a bulge of RNA silencing agent/target duplex. The intercalator can be an aromatic, *e.g.*, a polycyclic aromatic or heterocyclic aromatic compound. A polycyclic intercalator can have stacking capabilities, and can include systems with 2, 3, or 4 fused rings. The universal bases described herein can be included on a ligand. In one embodiment, the ligand can include a cleaving group that contributes to target gene inhibition by cleavage of the target nucleic acid. The cleaving group can be, for example, a bleomycin (*e.g.*, bleomycin-A5, bleomycin-A2, or bleomycin-B2), pyrene, phenanthroline (*e.g.*, O-phenanthroline), a polyamine, a tripeptide (*e.g.*, lys-tyr-lys tripeptide), or metal ion chelating group. The metal ion chelating group can include, *e.g.*, an Lu(III) or EU(III) macrocyclic complex, a Zn(II) 2,9-dimethylphenanthroline derivative, a Cu(II) terpyridine, or acridine, which can promote the selective cleavage of target RNA at the site of the bulge by free metal ions, such as Lu(III). In some embodiments, a peptide ligand can be tethered to a RNA silencing agent to promote cleavage of the target RNA, *e.g.*, at the bulge region. For example, 1,8-dimethyl-1,3,6,8,10,13-hexaazacyclotetradecane (cyclam) can be conjugated to a peptide (*e.g.*, by an amino acid derivative) to promote target RNA cleavage. A tethered ligand can be an aminoglycoside ligand, which can cause an RNA silencing agent to have improved hybridization properties or improved sequence specificity. Exemplary aminoglycosides include glycosylated polylysine, galactosylated polylysine, neomycin B, tobramycin, kanamycin A, and acridine conjugates of aminoglycosides, such as Neo-N-acridine, Neo-S-acridine, Neo-C-acridine, Tobra-N-acridine, and KanaA-N-acridine. Use of an acridine analog can increase sequence specificity. For example, neomycin B has a high affinity for RNA as compared to DNA, but low sequence-specificity. An acridine analog, neo-S-acridine has an increased affinity for the HIV Rev-response element (RRE). In some embodiments the guanidine analog (the guanidinoglycoside) of an aminoglycoside ligand is tethered to an RNA silencing agent. In a guanidinoglycoside, the amine group on the amino acid is exchanged for a guanidine group. Attachment of a guanidine analog can enhance cell permeability of an RNA silencing agent. A tethered ligand can be a poly-arginine

peptide, peptoid or peptidomimetic, which can enhance the cellular uptake of an oligonucleotide agent.

Preferred ligands are coupled, preferably covalently, either directly or indirectly via an intervening tether, to a ligand-conjugated carrier. In preferred embodiments, the ligand is attached to the carrier *via* an intervening tether. In preferred embodiments, a ligand alters the distribution, targeting or lifetime of an RNA silencing agent into which it is incorporated. In preferred embodiments a ligand provides an enhanced affinity for a selected target, *e.g.*, molecule, cell or cell type, compartment, *e.g.*, a cellular or organ compartment, tissue, organ or region of the body, as, *e.g.*, compared to a species absent such a ligand.

Preferred ligands can improve transport, hybridization, and specificity properties and may also improve nuclease resistance of the resultant natural or modified RNA silencing agent, or a polymeric molecule comprising any combination of monomers described herein and/or natural or modified ribonucleotides.

Ligands in general can include therapeutic modifiers, *e.g.*, for enhancing uptake; diagnostic compounds or reporter groups *e.g.*, for monitoring distribution; cross-linking agents; nuclease-resistance conferring moieties; and natural or unusual nucleobases. General examples include lipophiles, lipids, steroids (*e.g.*, uvaol, hecigenin, diosgenin), terpenes (*e.g.*, triterpenes, *e.g.*, sarsasapogenin, Friedelin, epifriedelanol derivatized lithocholic acid), vitamins (*e.g.*, folic acid, vitamin A, biotin, pyridoxal), carbohydrates, proteins, protein binding agents, integrin targeting molecules, polycationics, peptides, polyamines, and peptide mimics.

Ligands can include a naturally occurring substance, (*e.g.*, human serum albumin (HSA), low-density lipoprotein (LDL), or globulin); carbohydrate (*e.g.*, a dextran, pullulan, chitin, chitosan, inulin, cyclodextrin or hyaluronic acid); amino acid, or a lipid. The ligand may also be a recombinant or synthetic molecule, such as a synthetic polymer, *e.g.*, a synthetic polyamino acid. Examples of polyamino acids include polyamino acid is a polylysine (PLL), poly L-aspartic acid, poly L-glutamic acid, styrene-maleic acid anhydride copolymer, poly(L-lactide-co-glycolid) copolymer, divinyl ether-maleic anhydride copolymer, N-(2-hydroxypropyl)methacrylamide copolymer (HMPA), polyethylene glycol (PEG), polyvinyl alcohol (PVA), polyurethane, poly(2-ethylacrylic acid), N-isopropylacrylamide polymers, or

polyphosphazine. Example of polyamines include: polyethylenimine, polylysine (PLL), spermine, spermidine, polyamine, pseudopeptide-polyamine, peptidomimetic polyamine, dendrimer polyamine, arginine, amidine, protamine, cationic lipid, cationic porphyrin, quaternary salt of a polyamine, or an alpha helical peptide.

Ligands can also include targeting groups, *e.g.*, a cell or tissue targeting agent, *e.g.*, a lectin, glycoprotein, lipid or protein, *e.g.*, an antibody, that binds to a specified cell type such as a kidney cell. A targeting group can be a thyrotropin, melanotropin, lectin, glycoprotein, surfactant protein A, Mucin carbohydrate, multivalent lactose, multivalent galactose, N-acetyl-galactosamine, N-acetyl-glucosamine, multivalent mannose, multivalent fucose, glycosylated polyaminoacids, multivalent galactose, transferrin, bisphosphonate, polyglutamate, polyaspartate, a lipid, cholesterol, a steroid, bile acid, folate, vitamin B12, biotin, or an RGD peptide or RGD peptide mimetic.

Other examples of ligands include dyes, intercalating agents (*e.g.* acridines and substituted acridines), cross-linkers (*e.g.* psoralene, mitomycin C), porphyrins (TPPC4, texaphyrin, Sapphyrin), polycyclic aromatic hydrocarbons (*e.g.*, phenazine, dihydrophenazine, phenanthroline, pyrenes), lys-tyr-lys tripeptide, aminoglycosides, guanidium aminoglycosides, artificial endonucleases (*e.g.* EDTA), lipophilic molecules, *e.g.* cholesterol (and thio analogs thereof), cholic acid, cholanic acid, lithocholic acid, adamantane acetic acid, 1-pyrene butyric acid, dihydrotestosterone, glycerol (*e.g.*, esters (*e.g.*, mono, bis, or tris fatty acid esters, *e.g.*, C₁₀, C₁₁, C₁₂, C₁₃, C₁₄, C₁₅, C₁₆, C₁₇, C₁₈, C₁₉, or C₂₀ fatty acids) and ethers thereof, *e.g.*, C₁₀, C₁₁, C₁₂, C₁₃, C₁₄, C₁₅, C₁₆, C₁₇, C₁₈, C₁₉, or C₂₀ alkyl; *e.g.*, 1,3-bis-O(hexadecyl)glycerol, 1,3-bis-O(octadecyl)glycerol), geranyloxyhexyl group, hexadecylglycerol, borneol, menthol, 1,3-propanediol, heptadecyl group, palmitic acid, stearic acid (*e.g.*, glyceryl distearate), oleic acid, myristic acid, O3-(oleoyl)lithocholic acid, O3-(oleoyl)cholenic acid, dimethoxytrityl, or phenoxazine) and peptide conjugates (*e.g.*, antennapedia peptide, Tat peptide), alkylating agents, phosphate, amino, mercapto, PEG (*e.g.*, PEG-40K), MPEG, [MPEG]₂, polyamino, alkyl, substituted alkyl, radiolabeled markers, enzymes, haptens (*e.g.* biotin), transport/absorption facilitators (*e.g.*, aspirin, naproxen, vitamin E, folic acid), synthetic ribonucleases (*e.g.*, imidazole, bisimidazole, histamine, imidazole clusters, acridine-imidazole conjugates, Eu³⁺ complexes of tetraazamacrocycles), dinitrophenyl, HRP, or AP.

Ligands can be proteins, *e.g.*, glycoproteins, or peptides, *e.g.*, molecules having a specific affinity for a co-ligand, or antibodies *e.g.*, an antibody, that binds to a specified cell type such as a cancer cell, endothelial cell, or bone cell. Ligands may also include hormones and hormone receptors. They can also include non-peptidic species, such as lipids, lectins, carbohydrates, vitamins, cofactors, multivalent lactose, multivalent galactose, N-acetyl-galactosamine, N-acetyl-gulucosamine multivalent mannose, or multivalent fucose. The ligand can be, for example, a lipopolysaccharide, an activator of p38 MAP kinase, or an activator of NF- κ B.

The ligand can be a substance, *e.g.*, a drug, which can increase the uptake of the RNA silencing agent into the cell, for example, by disrupting the cell's cytoskeleton, *e.g.*, by disrupting the cell's microtubules, microfilaments, and/or intermediate filaments. The drug can be, for example, taxon, vincristine, vinblastine, cytochalasin, nocodazole, japlakinolide, latrunculin A, phalloidin, swinholide A, indanocine, or myoservin. The ligand can increase the uptake of the RNA silencing agent into the cell by activating an inflammatory response, for example. Exemplary ligands that would have such an effect include tumor necrosis factor alpha (TNFalpha), interleukin-1 beta, or gamma interferon. In one aspect, the ligand is a lipid or lipid-based molecule. Such a lipid or lipid-based molecule preferably binds a serum protein, *e.g.*, human serum albumin (HSA). An HSA binding ligand allows for distribution of the conjugate to a target tissue, *e.g.*, a non-kidney target tissue of the body. For example, the target tissue can be the liver, including parenchymal cells of the liver. Other molecules that can bind HSA can also be used as ligands. For example, neproxin or aspirin can be used. A lipid or lipid-based ligand can (a) increase resistance to degradation of the conjugate, (b) increase targeting or transport into a target cell or cell membrane, and/or (c) can be used to adjust binding to a serum protein, *e.g.*, HSA. A lipid based ligand can be used to modulate, *e.g.*, control the binding of the conjugate to a target tissue. For example, a lipid or lipid-based ligand that binds to HSA more strongly will be less likely to be targeted to the kidney and therefore less likely to be cleared from the body. A lipid or lipid-based ligand that binds to HSA less strongly can be used to target the conjugate to the kidney. In a preferred embodiment, the lipid based ligand binds HSA. A lipid-based ligand can bind HSA with a sufficient affinity such that the conjugate will be preferably distributed to a non-kidney tissue. However, it is preferred that the affinity not be so strong that the HSA-ligand binding cannot be reversed. In another preferred

embodiment, the lipid based ligand binds HSA weakly or not at all, such that the conjugate will be preferably distributed to the kidney. Other moieties that target to kidney cells can also be used in place of or in addition to the lipid based ligand.

In another aspect, the ligand is a moiety, *e.g.*, a vitamin, which is taken up by a target cell, *e.g.*, a proliferating cell. These are particularly useful for treating disorders characterized by unwanted cell proliferation, *e.g.*, of the malignant or non-malignant type, *e.g.*, cancer cells. Exemplary vitamins include vitamin A, E, and K. Other exemplary vitamins include are B vitamin, *e.g.*, folic acid, B12, riboflavin, biotin, pyridoxal or other vitamins or nutrients taken up by cancer cells. Also included are HSA and low density lipoprotein (LDL).

In another aspect, the ligand is a cell-permeation agent, preferably a helical cell-permeation agent. Preferably, the agent is amphipathic. An exemplary agent is a peptide such as tat or antennopodia. If the agent is a peptide, it can be modified, including a peptidylmimetic, invertomers, non-peptide or pseudo-peptide linkages, and use of D-amino acids. The helical agent is preferably an alpha-helical agent, which preferably has a lipophilic and a lipophobic phase.

The ligand can be a peptide or peptidomimetic. A peptidomimetic (also referred to herein as an oligopeptidomimetic) is a molecule capable of folding into a defined three-dimensional structure similar to a natural peptide. The attachment of peptide and peptidomimetics to oligonucleotide agents can affect pharmacokinetic distribution of the RNA silencing agent, such as by enhancing cellular recognition and absorption. The peptide or peptidomimetic moiety can be about 5-50 amino acids long, *e.g.*, about 5, 10, 15, 20, 25, 30, 35, 40, 45, or 50 amino acids long. A peptide or peptidomimetic can be, for example, a cell permeation peptide, cationic peptide, amphipathic peptide, or hydrophobic peptide (*e.g.*, consisting primarily of Tyr, Trp or Phe). The peptide moiety can be a dendrimer peptide, constrained peptide or crosslinked peptide. The peptide moiety can be an L-peptide or D-peptide. In another alternative, the peptide moiety can include a hydrophobic membrane translocation sequence (MTS). A peptide or peptidomimetic can be encoded by a random sequence of DNA, such as a peptide identified from a phage-display library, or one-bead-one-compound (OBOC) combinatorial library (Lam *et al.*, *Nature* 354:82-84, 1991). Preferably the peptide or peptidomimetic tethered to an RNA silencing agent via an incorporated monomer unit is a cell targeting peptide such as an arginine-glycine-aspartic acid (RGD)-peptide, or

RGD mimic. A peptide moiety can range in length from about 5 amino acids to about 40 amino acids. The peptide moieties can have a structural modification, such as to increase stability or direct conformational properties. Any of the structural modifications described below can be utilized.

IV. Methods of Introducing Nucleic Acids, Vectors, and Host Cells

RNA silencing agents of the invention may be directly introduced into the cell (*e.g.*, a neural cell) (*i.e.*, intracellularly); or introduced extracellularly into a cavity, interstitial space, into the circulation of an organism, introduced orally, or may be introduced by bathing a cell or organism in a solution containing the nucleic acid. Vascular or extravascular circulation, the blood or lymph system, and the cerebrospinal fluid are sites where the nucleic acid may be introduced.

The RNA silencing agents of the invention can be introduced using nucleic acid delivery methods known in art including injection of a solution containing the nucleic acid, bombardment by particles covered by the nucleic acid, soaking the cell or organism in a solution of the nucleic acid, or electroporation of cell membranes in the presence of the nucleic acid. Other methods known in the art for introducing nucleic acids to cells may be used, such as lipid-mediated carrier transport, chemical-mediated transport, and cationic liposome transfection such as calcium phosphate, and the like. The nucleic acid may be introduced along with other components that perform one or more of the following activities: enhance nucleic acid uptake by the cell or other-wise increase inhibition of the target gene.

The cell having the target gene may be from the germ line or somatic, totipotent or pluripotent, dividing or non-dividing, parenchyma or epithelium, immortalized or transformed, or the like. The cell may be a stem cell or a differentiated cell. Cell types that are differentiated include adipocytes, fibroblasts, myocytes, cardiomyocytes, endothelium, neurons, glia, blood cells, megakaryocytes, lymphocytes, macrophages, neutrophils, eosinophils, basophils, mast cells, leukocytes, granulocytes, keratinocytes, chondrocytes, osteoblasts, osteoclasts, hepatocytes, and cells of the endocrine or exocrine glands.

Depending on the particular target gene and the dose of RNA silencing agent material delivered, this process may provide partial or complete loss of function for the

target gene. A reduction or loss of gene expression in at least 50%, 60%, 70%, 80%, 90%, 95% or 99% or more of targeted cells is exemplary. Inhibition of gene expression refers to the absence (or observable decrease) in the level of protein and/or mRNA product from a target gene. Specificity refers to the ability to inhibit the target gene without manifest effects on other genes of the cell. The consequences of inhibition can be confirmed by examination of the outward properties of the cell or organism (as presented below in the examples) or by biochemical techniques such as RNA solution hybridization, nuclease protection, Northern hybridization, reverse transcription, gene expression monitoring with a microarray, antibody binding, enzyme linked immunosorbent assay (ELISA), Western blotting, radioimmunoassay (RIA), other immunoassays, and fluorescence activated cell analysis (FACS).

For RNA-mediated inhibition in a cell line or whole organism, gene expression is conveniently assayed by use of a reporter or drug resistance gene whose protein product is easily assayed. Such reporter genes include acetohydroxyacid synthase (AHAS), alkaline phosphatase (AP), beta galactosidase (LacZ), beta glucuronidase (GUS), chloramphenicol acetyltransferase (CAT), green fluorescent protein (GFP), horseradish peroxidase (HRP), luciferase (Luc), nopaline synthase (NOS), octopine synthase (OCS), and derivatives thereof. Multiple selectable markers are available that confer resistance to ampicillin, bleomycin, chloramphenicol, gentamycin, hygromycin, kanamycin, lincomycin, methotrexate, phosphinothricin, puromycin, and tetracyclin. Depending on the assay, quantitation of the amount of gene expression allows one to determine a degree of inhibition which is greater than 10%, 33%, 50%, 90%, 95% or 99% as compared to a cell not treated according to the present invention. Lower doses of injected material and longer times after administration of RNA silencing agent may result in inhibition in a smaller fraction of cells (*e.g.*, at least 10%, 20%, 50%, 75%, 90%, or 95% of targeted cells). Quantitation of gene expression in a cell may show similar amounts of inhibition at the level of accumulation of target mRNA or translation of target protein. As an example, the efficiency of inhibition may be determined by assessing the amount of gene product in the cell; mRNA may be detected with a hybridization probe having a nucleotide sequence outside the region used for the inhibitory double-stranded RNA, or translated polypeptide may be detected with an antibody raised against the polypeptide sequence of that region.

The RNA silencing agent may be introduced in an amount which allows delivery of at least one copy per cell. Higher doses (*e.g.*, at least 5, 10, 100, 500 or 1000 copies per cell) of material may yield more effective inhibition; lower doses may also be useful for specific applications.

V. Pharmaceutical Compositions and Methods of Administration

The RNA silencing agents of the invention can be incorporated into pharmaceutical compositions. Such compositions typically include the nucleic acid molecule and a pharmaceutically acceptable carrier. As used herein the language "pharmaceutically acceptable carrier" includes saline, solvents, dispersion media, coatings, antibacterial and antifungal agents, isotonic and absorption delaying agents, and the like, compatible with pharmaceutical administration. Supplementary active compounds can also be incorporated into the compositions.

A pharmaceutical composition is formulated to be compatible with its intended route of administration. Examples of routes of administration include parenteral, *e.g.*, intravenous, intradermal, subcutaneous, oral (*e.g.*, inhalation), transdermal (topical), transmucosal, and rectal administration. Solutions or suspensions used for parenteral, intradermal, or subcutaneous application can include the following components: a sterile diluent such as water for injection, saline solution, fixed oils, polyethylene glycols, glycerine, propylene glycol or other synthetic solvents; antibacterial agents such as benzyl alcohol or methyl parabens; antioxidants such as ascorbic acid or sodium bisulfite; chelating agents such as ethylenediaminetetraacetic acid; buffers such as acetates, citrates or phosphates and agents for the adjustment of tonicity such as sodium chloride or dextrose. pH can be adjusted with acids or bases, such as hydrochloric acid or sodium hydroxide. The parenteral preparation can be enclosed in ampoules, disposable syringes or multiple dose vials made of glass or plastic.

Pharmaceutical compositions suitable for injectable use include sterile aqueous solutions (where water soluble) or dispersions and sterile powders for the extemporaneous preparation of sterile injectable solutions or dispersion. For intravenous administration, suitable carriers include physiological saline, bacteriostatic water, Cremophor EL™ (BASF, Parsippany, NJ) or phosphate buffered saline (PBS). In all cases, the composition must be sterile and should be fluid to the extent that easy syringability exists. It should be stable under the conditions of manufacture and storage

and must be preserved against the contaminating action of microorganisms such as bacteria and fungi. The carrier can be a solvent or dispersion medium containing, for example, water, ethanol, polyol (for example, glycerol, propylene glycol, and liquid polyethylene glycol, and the like), and suitable mixtures thereof. The proper fluidity can be maintained, for example, by the use of a coating such as lecithin, by the maintenance of the required particle size in the case of dispersion and by the use of surfactants. Prevention of the action of microorganisms can be achieved by various antibacterial and antifungal agents, for example, parabens, chlorobutanol, phenol, ascorbic acid, thimerosal, and the like. In many cases, it will be preferable to include isotonic agents, for example, sugars, polyalcohols such as manitol, sorbitol, sodium chloride in the composition. Prolonged absorption of the injectable compositions can be brought about by including in the composition an agent which delays absorption, for example, aluminum monostearate and gelatin.

Sterile injectable solutions can be prepared by incorporating the active compound in the required amount in an appropriate solvent with one or a combination of ingredients enumerated above, as required, followed by filtered sterilization. Generally, dispersions are prepared by incorporating the active compound into a sterile vehicle, which contains a basic dispersion medium and the required other ingredients from those enumerated above. In the case of sterile powders for the preparation of sterile injectable solutions, the preferred methods of preparation are vacuum drying and freeze-drying which yields a powder of the active ingredient plus any additional desired ingredient from a previously sterile-filtered solution thereof.

Oral compositions generally include an inert diluent or an edible carrier. For the purpose of oral therapeutic administration, the active compound can be incorporated with excipients and used in the form of tablets, troches, or capsules, *e.g.*, gelatin capsules. Oral compositions can also be prepared using a fluid carrier for use as a mouthwash. Pharmaceutically compatible binding agents, and/or adjuvant materials can be included as part of the composition. The tablets, pills, capsules, troches and the like can contain any of the following ingredients, or compounds of a similar nature: a binder such as microcrystalline cellulose, gum tragacanth or gelatin; an excipient such as starch or lactose, a disintegrating agent such as alginic acid, Primogel, or corn starch; a lubricant such as magnesium stearate or Sterotes; a glidant such as colloidal silicon

dioxide; a sweetening agent such as sucrose or saccharin; or a flavoring agent such as peppermint, methyl salicylate, or orange flavoring.

For administration by inhalation, the RNA silencing agent may be delivered in the form of an aerosol spray from pressured container or dispenser which contains a suitable propellant, *e.g.*, a gas such as carbon dioxide, or a nebulizer. Such methods include those described in U.S. Patent No. 6,468,798.

Systemic administration can also be by transmucosal or transdermal means. For transmucosal or transdermal administration, penetrants appropriate to the barrier to be permeated are used in the formulation. Such penetrants are generally known in the art, and include, for example, for transmucosal administration, detergents, bile salts, and fusidic acid derivatives. Transmucosal administration can be accomplished through the use of nasal sprays or suppositories. For transdermal administration, the active compounds are formulated into ointments, salves, gels, or creams as generally known in the art.

The RNA silencing agents can also be prepared in the form of suppositories (*e.g.*, with conventional suppository bases such as cocoa butter and other glycerides) or retention enemas for rectal delivery.

The RNA silencing agents can also be administered by transfection or infection using methods known in the art, including but not limited to the methods described in McCaffrey et al. (2002), *Nature*, 418(6893), 38-9 (hydrodynamic transfection); Xia et al. (2002), *Nature Biotechnol.*, 20(10), 1006-10 (viral-mediated delivery); or Putnam (1996), *Am. J. Health Syst. Pharm.* 53(2), 151-160, erratum at *Am. J. Health Syst. Pharm.* 53(3), 325 (1996).

The RNA silencing agents can also be administered by any method suitable for administration of nucleic acid agents, such as a DNA vaccine. These methods include gene guns, bio-injectors, and skin patches as well as needle-free methods such as the micro-particle DNA vaccine technology disclosed in U.S. Patent No. 6,194,389, and the mammalian transdermal needle-free vaccination with powder-form vaccine as disclosed in U.S. Patent No. 6,168,587. Additionally, intranasal delivery is possible, as described in, *inter alia*, Hamajima et al. (1998), *Clin. Immunol. Immunopathol.*, 88(2), 205-10. Liposomes (*e.g.*, as described in U.S. Patent No. 6,472,375) and microencapsulation can also be used. Biodegradable targetable microparticle delivery systems can also be used (*e.g.*, as described in U.S. Patent No. 6,471,996).

In one embodiment, the RNA silencing agents are prepared with carriers that will protect the compound against rapid elimination from the body, such as a controlled release formulation, including implants and microencapsulated delivery systems. Biodegradable, biocompatible polymers can be used, such as ethylene vinyl acetate, polyanhydrides, polyglycolic acid, collagen, polyorthoesters, and polylactic acid. Such formulations can be prepared using standard techniques. The materials can also be obtained commercially from Alza Corporation and Nova Pharmaceuticals, Inc. Liposomal suspensions (including liposomes targeted to infected cells with monoclonal antibodies to viral antigens) can also be used as pharmaceutically acceptable carriers. These can be prepared according to methods known to those skilled in the art, for example, as described in U.S. Patent No. 4,522,811.

Toxicity and therapeutic efficacy of RNA silencing agents can be determined by standard pharmaceutical procedures in cell cultures or experimental animals, e.g., for determining the LD₅₀ (the dose lethal to 50% of the population) and the ED₅₀ (the dose therapeutically effective in 50% of the population). The dose ratio between toxic and therapeutic effects is the therapeutic index and it can be expressed as the ratio LD₅₀/ED₅₀. RNA silencing agents which exhibit high therapeutic indices are preferred. While RNA silencing agents that exhibit toxic side effects may be used, care should be taken to design a delivery system that targets such compounds to the site of affected tissue in order to minimize potential damage to uninfected cells and, thereby, reduce side effects.

The data obtained from the cell culture assays and animal studies can be used in formulating a range of dosage for use in humans. The dosage of such RNA silencing agents lies preferably within a range of circulating concentrations that include the ED₅₀ with little or no toxicity. The dosage may vary within this range depending upon the dosage form employed and the route of administration utilized. For any RNA silencing agent used in the method of the invention, the therapeutically effective dose can be estimated initially from cell culture assays. A dose may be formulated in animal models to achieve a circulating plasma concentration range that includes the IC₅₀ (*i.e.*, the concentration of the test RNA silencing agent which achieves a half-maximal inhibition of symptoms) as determined in cell culture. Such information can be used to more accurately determine useful doses in humans. Levels in plasma may be measured, for example, by high performance liquid chromatography.

As defined herein, a therapeutically effective amount of a RNA silencing agent (*i.e.*, an effective dosage) depends on the RNA silencing agent selected. For instance, if a plasmid encoding shRNA is selected, single dose amounts in the range of approximately 1 :g to 1000 mg may be administered; in some embodiments, 10, 30, 100 or 1000 :g may be administered. In some embodiments, 1-5 g of the compositions can be administered. The compositions can be administered one from one or more times per day to one or more times per week; including once every other day. The skilled artisan will appreciate that certain factors may influence the dosage and timing required to effectively treat a subject, including but not limited to the severity of the disease or disorder, previous treatments, the general health and/or age of the subject, and other diseases present. Moreover, treatment of a subject with a therapeutically effective amount of a protein, polypeptide, or antibody can include a single treatment or, preferably, can include a series of treatments.

The nucleic acid molecules of the invention can be inserted into expression constructs, *e.g.*, viral vectors, retroviral vectors, expression cassettes, or plasmid viral vectors, *e.g.*, using methods known in the art, including but not limited to those described in Xia et al., (2002), *supra*. Expression constructs can be delivered to a subject by, for example, inhalation, orally, intravenous injection, local administration (see U.S. Patent 5,328,470) or by stereotactic injection (see *e.g.*, Chen et al. (1994), Proc. Natl. Acad. Sci. USA, 91, 3054-3057). The pharmaceutical preparation of the delivery vector can include the vector in an acceptable diluent, or can comprise a slow release matrix in which the delivery vehicle is imbedded. Alternatively, where the complete delivery vector can be produced intact from recombinant cells, *e.g.*, retroviral vectors, the pharmaceutical preparation can include one or more cells which produce the gene delivery system.

The nucleic acid molecules of the invention can also include small hairpin RNAs (shRNAs), and expression constructs engineered to express shRNAs. Transcription of shRNAs is initiated at a polymerase III (pol III) promoter, and is thought to be terminated at position 2 of a 4-5-thymine transcription termination site. Upon expression, shRNAs are thought to fold into a stem-loop structure with 3' UU-overhangs; subsequently, the ends of these shRNAs are processed, converting the shRNAs into siRNA-like molecules of about 21 nucleotides. Brummelkamp et al. (2002), Science, 296, 550-553; Lee et al, (2002). *supra*; Miyagishi and Taira (2002),

Nature Biotechnol., 20, 497-500; Paddison et al. (2002), *supra*; Paul (2002), *supra*; Sui (2002) *supra*; Yu et al. (2002), *supra*.

The expression constructs may be any construct suitable for use in the appropriate expression system and include, but are not limited to retroviral vectors, linear expression cassettes, plasmids and viral or virally-derived vectors, as known in the art. Such expression constructs may include one or more inducible promoters, RNA Pol III promoter systems such as U6 snRNA promoters or H1 RNA polymerase III promoters, or other promoters known in the art. The constructs can include one or both strands of the siRNA. Expression constructs expressing both strands can also include loop structures linking both strands, or each strand can be separately transcribed from separate promoters within the same construct. Each strand can also be transcribed from a separate expression construct, Tuschl (2002), *supra*.

In certain exemplary embodiments, a composition that includes an RNA silencing agent of the invention can be delivered to the nervous system of a subject by a variety of routes. Exemplary routes include intrathecal, parenchymal (*e.g.*, in the brain), nasal, and ocular delivery. The composition can also be delivered systemically, *e.g.*, by intravenous, subcutaneous or intramuscular injection, which is particularly useful for delivery of the RNA silencing agents to peripheral neurons. A preferred route of delivery is directly to the brain, *e.g.*, into the ventricles or the hypothalamus of the brain, or into the lateral or dorsal areas of the brain. The RNA silencing agents for neural cell delivery can be incorporated into pharmaceutical compositions suitable for administration. For example, compositions can include one or more species of an RNA silencing agent and a pharmaceutically acceptable carrier. The pharmaceutical compositions of the present invention may be administered in a number of ways depending upon whether local or systemic treatment is desired and upon the area to be treated. Administration may be topical (including ophthalmic, intranasal, transdermal), oral or parenteral. Parenteral administration includes intravenous drip, subcutaneous, intraperitoneal or intramuscular injection, intrathecal, or intraventricular (*e.g.*, intracerebroventricular) administration.

The route of delivery can be dependent on the disorder of the patient. For example, a subject diagnosed with HD can be administered an anti-htt RNA silencing agent of the invention directly into the brain (*e.g.*, into the globus pallidus or the corpus striatum of the basal ganglia, and near the medium spiny neurons of the corpus striatum). In addition to an RNA silencing agent of the invention, a patient can be administered a second therapy, *e.g.*, a palliative therapy and/or disease-specific therapy. The secondary therapy can be, for example, symptomatic, (*e.g.*, for alleviating symptoms), neuroprotective (*e.g.*, for slowing or halting disease progression), or restorative (*e.g.*, for reversing the disease process). For the treatment of HD, for example, symptomatic therapies can include the drugs haloperidol, carbamazepine, or valproate. Other therapies can include psychotherapy, physiotherapy, speech therapy, communicative and memory aids, social support services, and dietary advice.

An RNA silencing agent can be delivered to neural cells of the brain. Delivery methods that do not require passage of the composition across the blood-brain barrier can be utilized. For example, a pharmaceutical composition containing an RNA silencing agent can be delivered to the patient by injection directly into the area containing the disease-affected cells. For example, the pharmaceutical composition can be delivered by injection directly into the brain. The injection can be by stereotactic injection into a particular region of the brain (*e.g.*, the substantia nigra, cortex, hippocampus, striatum, or globus pallidus). The RNA silencing agent can be delivered into multiple regions of the central nervous system (*e.g.*, into multiple regions of the brain, and/or into the spinal cord). The RNA silencing agent can be delivered into diffuse regions of the brain (*e.g.*, diffuse delivery to the cortex of the brain).

In one embodiment, the RNA silencing agent can be delivered by way of a cannula or other delivery device having one end implanted in a tissue, *e.g.*, the brain, *e.g.*, the substantia nigra, cortex, hippocampus, striatum or globus pallidus of the brain. The cannula can be connected to a reservoir of RNA silencing agent. The flow or delivery can be mediated by a pump, *e.g.*, an osmotic pump or minipump, such as an Alzet pump (Durect, Cupertino, CA). In one embodiment, a pump and reservoir are implanted in an area distant from the tissue, *e.g.*, in the abdomen, and delivery is effected by a conduit leading from the pump or reservoir to the site of release. Devices

for delivery to the brain are described, for example, in U.S. Patent Nos. 6,093,180, and 5,814,014.

An RNA silencing agent of the invention can be further modified such that it is capable of traversing the blood brain barrier. For example, the RNA silencing agent can be conjugated to a molecule that enables the agent to traverse the barrier. Such modified RNA silencing agents can be administered by any desired method, such as by intraventricular or intramuscular injection, or by pulmonary delivery, for example.

An RNA silencing agent of the invention can be administered ocularly, such as to treat retinal disorder, *e.g.*, a retinopathy. For example, the pharmaceutical compositions can be applied to the surface of the eye or nearby tissue, *e.g.*, the inside of the eyelid. They can be applied topically, *e.g.*, by spraying, in drops, as an eyewash, or an ointment. Ointments or droppable liquids may be delivered by ocular delivery systems known in the art such as applicators or eye droppers. Such compositions can include mucomimetics such as hyaluronic acid, chondroitin sulfate, hydroxypropyl methylcellulose or poly(vinyl alcohol), preservatives such as sorbic acid, EDTA or benzylchromium chloride, and the usual quantities of diluents and/or carriers. The pharmaceutical composition can also be administered to the interior of the eye, and can be introduced by a needle or other delivery device which can introduce it to a selected area or structure. The composition containing the RNA silencing agent can also be applied via an ocular patch.

In general, an RNA silencing agent of the invention can be administered by any suitable method. As used herein, topical delivery can refer to the direct application of an RNA silencing agent to any surface of the body, including the eye, a mucous membrane, surfaces of a body cavity, or to any internal surface. Formulations for topical administration may include transdermal patches, ointments, lotions, creams, gels, drops, sprays, and liquids. Conventional pharmaceutical carriers, aqueous, powder or oily bases, thickeners and the like may be necessary or desirable. Topical administration can also be used as a means to selectively deliver the RNA silencing agent to the epidermis or dermis of a subject, or to specific strata thereof, or to an underlying tissue.

Compositions for intrathecal or intraventricular (*e.g.*, intracerebroventricular) administration may include sterile aqueous solutions which may also contain buffers, diluents and other suitable additives. Compositions for intrathecal or intraventricular administration preferably do not include a transfection reagent or an additional lipophilic moiety besides, for example, the lipophilic moiety attached to the RNA silencing agent.

Formulations for parenteral administration may include sterile aqueous solutions which may also contain buffers, diluents and other suitable additives. Intraventricular injection may be facilitated by an intraventricular catheter, for example, attached to a reservoir. For intravenous use, the total concentration of solutes should be controlled to render the preparation isotonic.

An RNA silencing agent of the invention can be administered to a subject by pulmonary delivery. Pulmonary delivery compositions can be delivered by inhalation of a dispersion so that the composition within the dispersion can reach the lung where it can be readily absorbed through the alveolar region directly into blood circulation. Pulmonary delivery can be effective both for systemic delivery and for localized delivery to treat diseases of the lungs. In one embodiment, an RNA silencing agent administered by pulmonary delivery has been modified such that it is capable of traversing the blood brain barrier.

Pulmonary delivery can be achieved by different approaches, including the use of nebulized, aerosolized, micellular and dry powder-based formulations. Delivery can be achieved with liquid nebulizers, aerosol-based inhalers, and dry powder dispersion devices. Metered-dose devices are preferred. One of the benefits of using an atomizer or inhaler is that the potential for contamination is minimized because the devices are self contained. Dry powder dispersion devices, for example, deliver drugs that may be readily formulated as dry powders. An RNA silencing agent composition may be stably stored as lyophilized or spray-dried powders by itself or in combination with suitable powder carriers. The delivery of a composition for inhalation can be mediated by a dosing timing element which can include a timer, a dose counter, time measuring device, or a time indicator which when incorporated into the device enables dose tracking, compliance monitoring, and/or dose triggering to a patient during administration of the aerosol medicament.

The types of pharmaceutical excipients that are useful as carriers include stabilizers such as human serum albumin (HSA), bulking agents such as carbohydrates, amino acids and polypeptides; pH adjusters or buffers; salts such as sodium chloride; and the like. These carriers may be in a crystalline or amorphous form or may be a mixture of the two.

Bulking agents that are particularly valuable include compatible carbohydrates, polypeptides, amino acids or combinations thereof. Suitable carbohydrates include monosaccharides such as galactose, D-mannose, sorbose, and the like; disaccharides, such as lactose, trehalose, and the like; cyclodextrins, such as 2-hydroxypropyl- β -cyclodextrin; and polysaccharides, such as raffinose, maltodextrins, dextrans, and the like; alditols, such as mannitol, xylitol, and the like. A preferred group of carbohydrates includes lactose, trehalose, raffinose maltodextrins, and mannitol. Suitable polypeptides include aspartame. Amino acids include alanine and glycine, with glycine being preferred.

Suitable pH adjusters or buffers include organic salts prepared from organic acids and bases, such as sodium citrate, sodium ascorbate, and the like; sodium citrate is preferred.

An RNA silencing agent of the invention can be administered by oral and nasal delivery. For example, drugs administered through these membranes have a rapid onset of action, provide therapeutic plasma levels, avoid first pass effect of hepatic metabolism, and avoid exposure of the drug to the hostile gastrointestinal (GI) environment. Additional advantages include easy access to the membrane sites so that the drug can be applied, localized and removed easily. In one embodiment, an RNA silencing agent administered by oral or nasal delivery has been modified to be capable of traversing the blood-brain barrier.

In one embodiment, unit doses or measured doses of a composition that include RNA silencing agents are dispensed by an implanted device. The device can include a sensor that monitors a parameter within a subject. For example, the device can include a pump, such as an osmotic pump and, optionally, associated electronics.

An RNA silencing agent can be packaged in a viral natural capsid or in a chemically or enzymatically produced artificial capsid or structure derived therefrom.

VI. Methods of Treatment

The present invention provides for both prophylactic and therapeutic methods of treating a subject at risk of (or susceptible to) of Huntington's disease.

"Treatment", or "treating" as used herein, is defined as the application or administration of a therapeutic agent (*e.g.*, a RNA agent or vector or transgene encoding same) to a patient, or application or administration of a therapeutic agent to an isolated tissue or cell line from a patient, who has the disease or disorder, a symptom of disease or disorder or a predisposition toward a disease or disorder, with the purpose to cure, heal, alleviate, relieve, alter, remedy, ameliorate, improve or affect the disease or disorder, the symptoms of the disease or disorder, or the predisposition toward disease.

In one aspect, the invention provides a method for preventing Huntington's disease in a subject, by administering to the subject a therapeutic agent (*e.g.*, a RNA silencing agent or vector or transgene encoding same). Subjects at risk for the disease can be identified by, for example, any or a combination of diagnostic or prognostic assays as described herein. Administration of a prophylactic agent can occur prior to the manifestation of symptoms characteristic of the disease or disorder, such that the disease or disorder is prevented or, alternatively, delayed in its progression.

Another aspect of the invention pertains to methods treating subjects therapeutically, *i.e.*, alter onset of symptoms of the disease or disorder. In an exemplary embodiment, the modulatory method of the invention involves contacting a cell expressing a gain-of-function mutant with a therapeutic agent (*e.g.*, a RNA silencing agent or vector or transgene encoding same) that is specific for a mutation within the gene, such that sequence specific interference with the gene is achieved. These methods can be performed *in vitro* (*e.g.*, by culturing the cell with the agent) or, alternatively, *in vivo* (*e.g.*, by administering the agent to a subject).

With regards to both prophylactic and therapeutic methods of treatment, such treatments may be specifically tailored or modified, based on knowledge obtained from the field of pharmacogenomics. "Pharmacogenomics", as used herein, refers to the application of genomics technologies such as gene sequencing, statistical genetics, and gene expression analysis to drugs in clinical development and on the market. More

specifically, the term refers the study of how a patient's genes determine his or her response to a drug (*e.g.*, a patient's "drug response phenotype", or "drug response genotype"). Thus, another aspect of the invention provides methods for tailoring an individual's prophylactic or therapeutic treatment with either the target gene molecules of the present invention or target gene modulators according to that individual's drug response genotype. Pharmacogenomics allows a clinician or physician to target prophylactic or therapeutic treatments to patients who will most benefit from the treatment and to avoid treatment of patients who will experience toxic drug-related side effects.

Therapeutic agents can be tested in an appropriate animal model. For example, an RNA silencing agent (or expression vector or transgene encoding same) as described herein can be used in an animal model to determine the efficacy, toxicity, or side effects of treatment with said agent. Alternatively, a therapeutic agent can be used in an animal model to determine the mechanism of action of such an agent. For example, an agent can be used in an animal model to determine the efficacy, toxicity, or side effects of treatment with such an agent. Alternatively, an agent can be used in an animal model to determine the mechanism of action of such an agent.

A pharmaceutical composition containing an RNA silencing agent of the invention can be administered to any patient diagnosed as having or at risk for developing a neurological disorder, such as HD. In one embodiment, the patient is diagnosed as having a neurological disorder, and the patient is otherwise in general good health. For example, the patient is not terminally ill, and the patient is likely to live at least 2, 3, 5, or 10 years or longer following diagnosis. The patient can be treated immediately following diagnosis, or treatment can be delayed until the patient is experiencing more debilitating symptoms, such as motor fluctuations and dyskinesia in PD patients. In another embodiment, the patient has not reached an advanced stage of the disease.

An RNA silencing agent modified for enhance uptake into neural cells can be administered at a unit dose less than about 1.4 mg per kg of bodyweight, or less than 10, 5, 2, 1, 0.5, 0.1, 0.05, 0.01, 0.005, 0.001, 0.0005, 0.0001, 0.00005 or 0.00001 mg per kg of bodyweight, and less than 200 nmole of RNA agent (*e.g.*, about 4.4×10^{16} copies) per kg of bodyweight, or less than 1500, 750, 300, 150, 75, 15, 7.5, 1.5, 0.75, 0.15, 0.075,

0.015, 0.0075, 0.0015, 0.00075, 0.00015 nmole of RNA silencing agent per kg of bodyweight. The unit dose, for example, can be administered by injection (*e.g.*, intravenous or intramuscular, intrathecally, or directly into the brain), an inhaled dose, or a topical application. Particularly preferred dosages are less than 2, 1, or 0.1 mg/kg of body weight.

Delivery of an RNA silencing agent directly to an organ (*e.g.*, directly to the brain) can be at a dosage on the order of about 0.00001 mg to about 3 mg per organ, or preferably about 0.0001-0.001 mg per organ, about 0.03- 3.0 mg per organ, about 0.1- 3.0 mg per eye or about 0.3-3.0 mg per organ. The dosage can be an amount effective to treat or prevent a neurological disease or disorder, *e.g.*, HD. In one embodiment, the unit dose is administered less frequently than once a day, *e.g.*, less than every 2, 4, 8 or 30 days. In another embodiment, the unit dose is not administered with a frequency (*e.g.*, not a regular frequency). For example, the unit dose may be administered a single time. In one embodiment, the effective dose is administered with other traditional therapeutic modalities.

In one embodiment, a subject is administered an initial dose, and one or more maintenance doses of an RNA silencing agent. The maintenance dose or doses are generally lower than the initial dose, *e.g.*, one-half less of the initial dose. A maintenance regimen can include treating the subject with a dose or doses ranging from 0.01 µg to 1.4 mg/kg of body weight per day, *e.g.*, 10, 1, 0.1, 0.01, 0.001, or 0.00001 mg per kg of bodyweight per day. The maintenance doses are preferably administered no more than once every 5, 10, or 30 days. Further, the treatment regimen may last for a period of time which will vary depending upon the nature of the particular disease, its severity and the overall condition of the patient. In preferred embodiments the dosage may be delivered no more than once per day, *e.g.*, no more than once per 24, 36, 48, or more hours, *e.g.*, no more than once every 5 or 8 days. Following treatment, the patient can be monitored for changes in his condition and for alleviation of the symptoms of the disease state. The dosage of the compound may either be increased in the event the patient does not respond significantly to current dosage levels, or the dose may be decreased if an alleviation of the symptoms of the disease state is observed, if the disease state has been ablated, or if undesired side-effects are observed.

The effective dose can be administered in a single dose or in two or more doses, as desired or considered appropriate under the specific circumstances. If desired to facilitate repeated or frequent infusions, implantation of a delivery device, *e.g.*, a pump, semi-permanent stent (*e.g.*, intravenous, intraperitoneal, intracisternal or intracapsular), or reservoir may be advisable. In one embodiment, a pharmaceutical composition includes a plurality of RNA silencing agent species. In another embodiment, the RNA silencing agent species has sequences that are non-overlapping and non-adjacent to another species with respect to a naturally occurring target sequence. In another embodiment, the plurality of RNA silencing agent species is specific for different naturally occurring target genes. In another embodiment, the RNA silencing agent is allele specific. In another embodiment, the plurality of RNA silencing agent species target two or more SNP alleles (*e.g.*, two, three, four, five, six, or more SNP alleles).

Following successful treatment, it may be desirable to have the patient undergo maintenance therapy to prevent the recurrence of the disease state, wherein the compound of the invention is administered in maintenance doses, ranging from 0.01 μ g to 100 g per kg of body weight (see US 6,107,094).

The concentration of the RNA silencing agent composition is an amount sufficient to be effective in treating or preventing a disorder or to regulate a physiological condition in humans. The concentration or amount of RNA silencing agent administered will depend on the parameters determined for the agent and the method of administration, *e.g.* nasal, buccal, or pulmonary. For example, nasal formulations tend to require much lower concentrations of some ingredients in order to avoid irritation or burning of the nasal passages. It is sometimes desirable to dilute an oral formulation up to 10-100 times in order to provide a suitable nasal formulation.

Certain factors may influence the dosage required to effectively treat a subject, including but not limited to the severity of the disease or disorder, previous treatments, the general health and/or age of the subject, and other diseases present. Moreover, treatment of a subject with a therapeutically effective amount of an RNA silencing agent can include a single treatment or, preferably, can include a series of treatments. It will also be appreciated that the effective dosage of an RNA silencing agent for treatment may increase or decrease over the course of a particular treatment. Changes in dosage

may result and become apparent from the results of diagnostic assays as described herein. For example, the subject can be monitored after administering an RNA silencing agent composition. Based on information from the monitoring, an additional amount of the RNA silencing agent composition can be administered.

Dosing is dependent on severity and responsiveness of the disease condition to be treated, with the course of treatment lasting from several days to several months, or until a cure is effected or a diminution of disease state is achieved. Optimal dosing schedules can be calculated from measurements of drug accumulation in the body of the patient. Persons of ordinary skill can easily determine optimum dosages, dosing methodologies and repetition rates. Optimum dosages may vary depending on the relative potency of individual compounds, and can generally be estimated based on EC50s found to be effective in *in vitro* and *in vivo* animal models. In some embodiments, the animal models include transgenic animals that express a human gene, *e.g.*, a gene that produces a target RNA, *e.g.*, an RNA expressed in a neural cell. The transgenic animal can be deficient for the corresponding endogenous RNA. In another embodiment, the composition for testing includes an RNA silencing agent that is complementary, at least in an internal region, to a sequence that is conserved between the target RNA in the animal model and the target RNA in a human.

VII. Kits

In certain other aspects, the invention provides kits that include a suitable container containing a pharmaceutical formulation of an RNA silencing agent, *e.g.*, a double-stranded RNA silencing agent, or sRNA agent, (*e.g.*, a precursor, *e.g.*, a larger RNA silencing agent which can be processed into a sRNA agent, or a DNA which encodes an RNA silencing agent, *e.g.*, a double-stranded RNA silencing agent, or sRNA agent, or precursor thereof). In certain embodiments the individual components of the pharmaceutical formulation may be provided in one container. Alternatively, it may be desirable to provide the components of the pharmaceutical formulation separately in two or more containers, *e.g.*, one container for an RNA silencing agent preparation, and at least another for a carrier compound. The kit may be packaged in a number of different configurations such as one or more containers in a single box. The different components can be combined, *e.g.*, according to instructions provided with the kit. The components

can be combined according to a method described herein, *e.g.*, to prepare and administer a pharmaceutical composition. The kit can also include a delivery device.

This invention is further illustrated by the following examples which should not be construed as limiting. The contents of all references, patents and published patent applications cited throughout this application are incorporated herein by reference.

EXAMPLES

Example I: SNP analysis in patients with Huntington's Disease (HD)

Single nucleotide polymorphism (SNP) sequencing analysis was performed to identify heterozygotic SNPs in patients with Huntington's Disease (HD). DNA samples were obtained from brain repositories in Charlestown, MA, USA and New York City, NY and a DNA repository in Ulm, Germany.

In one study, a total of 195 subjects (HD and control) were examined. Twenty-one SNP sites were identified; eighteen were reported in SNPPER and three have not previously been reported. In this population, four reported SNP sites were not confirmed. 149/195 subjects (76%) contained SNP heterozygosity. The USA and German populations had 76% heterozygosity at SNP sites. HD patients and controls had the same frequency of SNP heterozygosity. Six SNP sites had an allelic frequency >30%.

In a second study, a total of twelve candidate polymorphic sites were sequenced in more than 107 HD brain specimens. Fifty-five percent of HD patients in the USA specimens (26/47; USA) and 57% of controls (50/88; USA) were heterozygous at one or more SNP site. Thirty-eight percent of HD patients in the German collection (23/60; Germany) were heterozygous at one or more polymorphic site. Forty-three of the total 107 HD patients had clusters of heterozygous polymorphic sites, with heterozygosity at five or eight SNP sites.

Example II: Testing of RNA silencing agents (e.g., siRNAs) against htt SNPs

Two SNPs (at genomic positions RS 363125 and RS 362331) were selected to test RNAi discrimination in an *in vitro* reporter assay.

SiRNAs targeting each SNP were designed as described *supra*. siRNA strands were annealed (Elbashir et al., 2001a).

Target RNAs may be prepared as follows. Target RNAs are transcribed with recombinant, histidine-tagged, T7 RNA polymerase from PCR products as described (Nykänen et al., 2001; Hutvágner et al., 2002). PCR templates for htt sense and anti-sense target RNA preparation are generated by amplifying plasmid template encoding htt cDNA using primer pairs. Each primer pair consists of forward and reverse primers which are complementary to regions that are immediately upstream and downstream, respectively, of one of the two alleles of a heterozygous SNP site.

Each siRNA was tested using a cell-based *in vitro* luciferase reporter assay. *In vitro* RNAi reactions and analysis may also be carried out in *Drosophila* embryo lysates as previously described (Tuschl et al., 1999; Zamore et al., 2000; Haley et al., 2003). Target RNAs are used at ~ 5 nM concentration so that reactions are mainly under single-turnover conditions. Target cleavage under these conditions is proportionate to siRNA concentration.

In RS363125 (predicted high discriminator), 80% knockdown was achieved at 0.5 nM siRNA for a siRNA having a guide strand perfectly complementary to a target mRNA containing the polymorphic adenosine ("A") SNP allele of the target SNP site from *huntingtin*, but containing a mismatch with a non-target mRNA containing the corresponding cytosine ("C") SNP allele of the heterozygous RS363125 SNP site (Figure 4A). A single nucleotide mismatch effectively discriminated between both alleles of this SNP as the siRNA did not achieve the same level of knockdown of non-target ("mismatched") mRNA, even at up to 20 nM siRNA (Figure 4B). In contrast, an siRNA having perfect complementarity to the "C" SNP allele and a mismatch with the "A" SNP allele of the RS363125 SNP site (Figure 5A) was effective in achieving discriminatory RNAi in favor of the "C" SNP allele (Figure 5B).

Similar findings applied to the RS362331 polymorphism. The sequences of the 21-mer siRNAs targeting each SNP allele ("C" or "U") of the RS362331 polymorphism are depicted in Figure 2A (SEQ ID NO: 3 sense; SEQ ID NO: 4 anti-sense) and Figure 3A (SEQ ID NO: 7 sense; SEQ ID NO: 8 anti-sense). Each siRNA is perfectly complementary to the targeted SNP allele, but contains a mismatch with the non-targeted SNP allele at position 10 of the guide strand (P10).

Figure 2B shows the efficacy of a siRNA directed against the SNP allele of RS362331 having a cytosine ("C") at the heterozygous SNP site (SEQ ID NO:5; match). The data clearly demonstrate that the siRNA directs cleavage of the matched

target SNP allele to a greater degree than observed for the mismatched target SNP allele (SEQ ID NO:6; mismatch). Similarly, Figure 3B, shows the efficacy of siRNA against the SNP allele of RS362331 which has a uridine ("U") at the heterozygotic SNP site (SEQ ID NO: 6; match). Greater than 50% knockdown of the targeted SNP allele is achieved at 0.5nM siRNA, while the non-targeted "C" SNP allele (SEQ ID NO:5; mismatch) is relatively unaffected.

The efficacy of siRNA directed against the homozygous C allele (SEQ ID NO: 17) in the 3'UTR of Htt gene was also tested. siRNAs having a guide strand with perfect complementary to the C allele (Figure 6A) or containing a single U:C mismatch with the C allele at position 10 (P10) of the guide strand (Figure 6B) were designed. Transfection was performed at an siRNA concentration of 20nM in HEK cells homozygous for the C allele. Transfection efficiency was approximately 70%. As depicted in Figure 6C and Figure 6D, matched siRNAs were much more effective than mismatched siRNAs in achieving knockdown of both Htt target mRNA and protein levels, respectively. This pattern of gene silencing was consistently observed at lower siRNA concentrations (5nM and 10nM) as well (see Figure 7).

These results indicate that siRNAs can preferentially silence one of the two SNP alleles differing at the polymorphic site and that heterozygous SNP sites in *huntingtin* are attractive targets for therapeutic siRNAs.

Example III: Effect of RNA silencing agents (e.g., siRNAs) on Mutant htt Protein Expression

Western blotting may also be employed to test the ability of siRNAs to down-regulate endogenous Htt protein in cultured cells using any of the techniques described *infra*.

RNA silencing activity of SNP-specific siRNAs can be tested HeLa cells. HeLa cells are maintained at 37°C in Dulbecco's modified Eagle's medium (DMEM, Invitrogen) supplemented with 10% fetal bovine serum (FBS), 100 unit/ml penicillin and 100 µg/ml streptomycin (Invitrogen). Cells are regularly passaged at sub-confluence and plated at 70% confluency 16 hours before transfection. Lipofectamine™ (Invitrogen)-mediated transient transfection of siRNAs are performed in duplicate 6-well plates (Falcon) as described for adherent cell lines by the manufacturer. A standard transfection mixture containing 100-150 nM siRNA and 9-10 µl Lipofectamine™ in 1

ml serum-reduced OPTI-MEM® (Invitrogen) are added to each well. Cells are incubated in transfection mixture at 37°C for 6 hours and further cultured in antibiotic-free DMEM. For Western blot analysis at various time intervals, the transfected cells are harvested, washed twice with phosphate buffered saline (PBS, Invitrogen), flash frozen in liquid nitrogen, and stored at -80°C for analysis.

Western blot analysis is performed as follows. Cells treated with siRNA are harvested as described above and lysed in ice-cold reporter lysis buffer (Promega) containing protease inhibitor (complete, EDTA-free, 1 tablet/10 ml buffer, Roche Molecular Biochemicals). After clearing the resulting lysates by centrifugation, protein in clear lysates is quantified by Dc protein assay kit (Bio-Rad). Proteins in 60 µg of total cell lysate are resolved by 10% SDS-PAGE, transferred onto a polyvinylidene difluoride membrane (PVDF, Bio-Rad), and immuno-blotted with antibodies against CD80 (Santa Cruz). Protein content is visualized with a BM Chemiluminescence Blotting Kit (Roche Molecular Biochemicals). The blots are exposed to x-ray film (Kodak MR-1) for various times (30 s to 5 min).

siRNAs against polymorphic regions in the htt mRNA can likewise be tested in cells transfected with human htt cDNA or in cells transfected with htt reporter constructs. Lipofectamine™ (Invitrogen)-mediated transient cotransfections of cDNAs or reporter plasmids and siRNAs are performed as described *supra*. To test the ability of siRNAs to target htt reported constructs, RNAi is used to inhibit GFP-htt expression in cultured human HeLa cell lines. Briefly, HeLa cells are transfected with GFP-htt siRNA duplex, targeting the GFP-htt mRNA sequence. To analyze RNAi effects against GFP-htt, lysates are prepared from siRNA duplex-treated cells at various times after transfection. Western blot experiments are carried out as described *supra*.

Inhibition of htt expression can be assessed in cultured neuronal cells as well. Exemplary cells include PC12 (Scheitzer et al., Thompson et al.) and NT3293 (Tagle et al.) cell lines as previously described. Additional exemplary cells include stably-transfected cells, *e.g.* neuronal cells or neuronally-derived cells. PC12 cell lines expressing exon 1 of the human huntingtin gene (Htt) can be used although expression of exon 1 reduces cell survival. GFP-Htt PC12 cells having an inducible GFP-Htt gene can also be used to test or validate siRNA efficacy.

Example IV: Htt siRNA delivery in an *in vivo* setting

R6/2 mice models (expressing the R6/2 human htt cDNA product) are an accepted animal model to study the effectiveness of siRNA delivery in an *in vivo* setting. Genetically engineered R6/2 mice may be used to test the effectiveness of siRNA at the 5' terminus of huntingtin mRNA. Htt siRNA are injected into the striatum of R6/2 mice through an Alzet pump. Mice are treated for 14 days with the siRNA/Alzet pump delivery system.

Equivalents

Those skilled in the art will recognize, or be able to ascertain using no more than routine experimentation, many equivalents to the specific embodiments of the invention described herein. Such equivalents are intended to be encompassed by the following claims.

WHAT IS CLAIMED IS:

1. A method of silencing a target mRNA encoding a mutant huntingtin (htt) protein in a cell, comprising contacting the cell with effective amount of a RNA silencing agent targeting a heterozygous single nucleotide polymorphism (SNP) within the target mRNA, such that RNA silencing of said mRNA occurs, wherein the SNP has an allelic frequency of at least 35% in a sample population.
2. The method of claim 1, wherein the SNP is present at genomic site RS362331.
3. The method of claim 3, wherein the target mRNA comprises the sequence set forth as SEQ ID NO: 5.
4. The method of claim 3, wherein the target mRNA comprises the sequence set forth as SEQ ID NO: 6.
5. The method of claim 3, wherein the RNA silencing agent is capable of inducing discriminatory RNA silencing.
6. The method of claim 3, wherein the antisense strand of said RNA silencing agent is complementary to the SNP and wherein said RNA silencing agent is capable of substantially silencing the mutant huntingtin protein without substantially silencing the corresponding wild-type huntingtin protein.
7. The method of claim 3, wherein the sample population is of Western European origin.
8. A method of silencing a target mRNA encoding a mutant huntingtin (htt) protein in a cell, comprising contacting the cell with effective amount of a siRNA targeting a heterozygous single nucleotide polymorphism (SNP) within the target mRNA, such that RNA silencing of said mRNA occurs, wherein the siRNA is selected from the group consisting of :
 - a. an siRNA comprising (i) a sense strand comprising the sequence set forth as SEQ ID NO: 3 or a variant thereof; and (ii) an antisense strand comprising the sequence set forth as SEQ ID NO: 4 or a variant thereof,

- said variant comprising at least one nucleotide analog or backbone modification;
- b. an siRNA comprising (i) a sense strand comprising the sequence set forth as SEQ ID NO: 7 or a variant thereof; and (ii) an antisense strand comprising the sequence set forth as SEQ ID NO: 8 or a variant thereof, said variant comprising at least one nucleotide analog or backbone modification,
 - c. an siRNA comprising (i) a sense strand comprising the sequence set forth as SEQ ID NO: 9 or a variant thereof; and (ii) an antisense strand comprising the sequence set forth as SEQ ID NO: 10 or a variant thereof, said variant comprising at least one nucleotide analog or backbone modification, and
 - d. an siRNA comprising (i) a sense strand comprising the sequence set forth as SEQ ID NO: 13 or a variant thereof; and (ii) an antisense strand comprising the sequence set forth as SEQ ID NO: 14 or a variant thereof, said variant comprising at least one nucleotide analog or backbone modification.
9. The method of any of claim 7, wherein the siRNA comprises a lipophilic moiety.
10. The method of claim 9, wherein the lipophilic moiety is a cholesterol moiety.
11. An RNA silencing agent comprising an antisense strand comprising about 16-25 nucleotides homologous to a region of an mRNA encoding a mutant huntingtin (htt) protein, said region comprising a heterozygous single nucleotide polymorphism (SNP) having an allelic frequency of at least 35% in a sample population, wherein the RNA silencing agent is capable of directing RNA silencing of said mRNA.
12. The RNA silencing agent of claim 11, wherein the SNP is present at genomic site RS362331.

13. The RNA silencing agent of claim 11, wherein the target mRNA comprises the sequence set forth as SEQ ID NO: 5.
14. The RNA silencing agent of claim 11, wherein the target mRNA comprises the sequence set forth as SEQ ID NO: 6.
15. The RNA silencing agent of claim 11, which is capable of inducing discriminatory RNA silencing.
16. The RNA silencing agent of claim 11, which is capable of substantially silencing the mutant huntingtin protein without substantially silencing the corresponding wild-type huntingtin protein.
17. The RNA silencing agent of claim 11, wherein the RNA silencing agent is an siRNA.
18. The siRNA of claim 17, comprising a sense strand having a nucleotide sequence identical to the sequence of the SNP.
19. The siRNA of claim 11, wherein the sense strand of the siRNA is identical to the polymorphism at a nucleotide position that is 10 nucleotides from the 5' end of the sense strand.
20. The siRNA of claim 11, wherein the sense strand of the siRNA is identical to the polymorphism at a nucleotide position that is 16 nucleotides from the 5' end of the sense strand.
21. An siRNA is selected from the group consisting of :
 - a. an siRNA comprising (i) a sense strand comprising the sequence set forth as SEQ ID NO: 3 or a variant thereof; and (ii) an antisense strand comprising the sequence set forth as SEQ ID NO: 4 or a variant thereof, said variant comprising at least one nucleotide analog or backbone modification;
 - b. an siRNA comprising (i) a sense strand comprising the sequence set forth as SEQ ID NO: 7 or a variant thereof; and (ii) an antisense strand

comprising the sequence set forth as SEQ ID NO: 8 or a variant thereof, said variant comprising at least one nucleotide analog or backbone modification,

- c. an siRNA comprising (i) a sense strand comprising the sequence set forth as SEQ ID NO: 9 or a variant thereof; and (ii) an antisense strand comprising the sequence set forth as SEQ ID NO: 10 or a variant thereof, said variant comprising at least one nucleotide analog or backbone modification, and
- d. an siRNA comprising (i) a sense strand comprising the sequence set forth as SEQ ID NO: 13 or a variant thereof; and (ii) an antisense strand comprising the sequence set forth as SEQ ID NO: 14 or a variant thereof, said variant comprising at least one nucleotide analog or backbone modification.

22. The siRNA of claim 21, wherein the siRNA comprises a lipophilic moiety.

23. The siRNA of claim 22, wherein the lipophilic moiety is a cholesterol moiety.

Figure 1

SNP site (refSNP)	Location	Allelic Frequency	SNP alleles (gene)	SNP alleles (mRNA)
RS362331	Exon 50	42%	C/T	C/U
RS4690077	Exon 48	34.9%	A/G	A/G
RS363125	Exon 39	15.4%	A/C	A/C
47bp into	Exon 25	11.2%	G/A	G/A
RS363075	Exon 20	11.2%	G/A	G/A
RS362268	3'UTR	33.9%	C/G	C/G
RS362267	3'UTR	33.3%	C/T	C/U
RS362307	3'UTR	32.8%	C/T	C/U
RS362306	3'UTR	33.3%	A/G	A/G
RS362305	3'UTR	16.4%	C/T	C/U
RS362304	3'UTR	17.9%	C/A	C/A
RS362303	3'UTR	15.8%	C/T	C/U

Figure 2

A.

5' UCCCUCAUC ACUGUGUAAC 3' (SEQ ID NO: 3)
3' CGAGGGAGUAG UGACACAG5' (SEQ ID NO: 4)
5' CCACGCCUGCUCCCUCAUC ACUGUGGCACUUAUCCUG 3' C match (SEQ ID NO: 5)
5' CCACGCCUGCUCCCUCAUC ACUGUGGCACUUAUCCUG 3' C mismatch (SEQ ID NO: 6)

B. "C" Match / "U" Mismatch




Figure 3





Figure 4

A.

5' GGGACAGUAaUUCAACGGUC 3' (SEQ ID NO:9)
3' ACCCUGUCAUUAAGUUGCGA 5' (SEQ ID NO:10)

5' UAAAGAGAUGGGACAGUAaUUCAACGCUAGAAGAACA 3' (SEQ ID NO:11)
5' UAAAGAGAUGGGACAGUAaUUCAACGCUAGAAGAACA 3' (SEQ ID NO:12)

RS363125

"A" Match / "C" Mismatch

B.




Figure 5

A.

5' GGGACAGUAGUUCAACGCGUC 3' (SEQ ID NO:13)
 3' ACCCCUGUCAUGAAGUUGCGA 5' (SEQ ID NO:14)

5' UAGAGAUUGGGACAGUACUUCAACGCUAGAAGACA 3' (SEQ ID NO:12)
 5' UAGAGAUUGGGACAGUAAUUCACGCUAGAAGACA 3' (SEQ ID NO:11)

RS363125

"C" Match / "A" Mismatch

B.




Figure 6

A.

Match siRNA


5'-ugccauggccugugcuggucc-3' SEQ ID NO:15
 3'-cuacgguaaccgGacacgaccc-5' SEQ ID NO:16
 5'-cagatgcccatggcCgtgctggggccag-3' SEQ ID NO:17

B.

Mismatch siRNA

5'-ugccauggccugugcuggucc-3' SEQ ID NO:18
 3'-cuacgguaaccgGacacgaccc-5' SEQ ID NO:19
 5'-cagatgcccatggcCgtgctggggccag-3' SEQ ID NO:17

C.



D.




Figure 7